



Article Type: Research Article

Received: 21/06/2020

Published: 07/07/2020

DOI: 10.46718/JBGSR.2020.02.000058

Biodegradation of Polycyclic Aromatic Hydrocarbon Mixtures by *Rhodococcus Pyridinivorans* FF2 and *Pseudomonas Aeruginosa* F4b Isolated from Sediments of Lagos Lagoon, Nigeria

Chioma C Obi¹, Sunday A Adebuseye, Olukayode O Amund², Esther E Ugoji² and William J Hickey³¹Department of Biological Sciences, Bells University of Technology Ota, Ogun State, Nigeria²Department of Microbiology, University of Lagos, Akoka, Yaba, Lagos, Nigeria³Department of Soil Science, University of Wisconsin-Madison, Madison, WI, USA

*Corresponding author: William J Hickey, Department of Soil Science, University of Wisconsin-Madison, Madison, WI, USA

Abstract

Polycyclic aromatic hydrocarbons (PAH) are widespread environmental pollutants that need urgent attention because of their toxicity. Development of microbial inoculants for PAH bioremediation is a potential avenue by which the environmental hazards posed by PAH can be addressed. The goal of the study was to determine if using PAH mixtures, rather than single PAH, as enrichment substrates would yield isolates that have superior PAH degradation abilities and/or are retrieve novel taxa from the PAH-contaminated sediments of Lagos Lagoon, Nigeria. The use of a quaternary mixture of PAH (pyrene, fluoranthene, fluorene, benz[a]anthracene) was successful in yielding two isolates, *Pseudomonas aeruginosa* strain F4b and *Rhodococcus pyridinivorans* strain FF2 with capabilities to grow on multiple PAH, and thus potentially useful in bioremediation. In addition to the PAH degraded both isolates could grow on a wide range of other hydrocarbon substrates. The isolates of *P. aeruginosa* and *R. pyridinivorans* were identified as possessing PAH ring-hydroxylating dioxygenases of the nahAC and narAa genotypes, respectively. The present study extends our knowledge of PAH biodegradation by *P. aeruginosa* and is the first report of PAH biodegradation by *R. pyridinivorans*. The capability of the *R. pyridinivorans* isolate to effectively degrade a highly toxic PAH, benz[a]anthracene, has particular importance for use in bioremediation.

Introduction

In Nigeria, oil exploration and indiscriminate discharge of municipal and industrial effluents has resulted in a tremendous increase in environmental pollution by many organic compounds including polycyclic aromatic hydrocarbons (PAHs). These compounds, especially the high molecular weight PAHs (HMW-PAHs, composed of four or more rings) are suspected carcinogens and are very hazardous to humans when ingested, inhaled or through other direct contact [1,2]. The concentration of PAHs in the environment is usually directly proportional to the type of human activities inherent in such environment [3]. Thus, the major source of PAHs in coastal sediments is usually through anthropogenic sources such as atmospheric

deposition, urban runoffs, municipal and industrial effluents agricultural wastes and oil spillages [4,5].

PAHs may undergo adsorption, volatilization, photolysis, and chemical degradation but microbial degradation is recognized as the main pathway of PAHs removal from a contaminated site [6,7]. PAH degradation depends on the environmental conditions, number and type of the microorganisms, nature and chemical structure of the chemical compound being degraded [8]. Microbial degradation of PAHs in the environment can occur through metabolism or co-metabolism [9]. Bioremediation of hydrocarbon-contaminated sites relies on either the activity of microorganisms already present at such site or the

addition of selected microorganisms that have the suitable enzymatic systems with desired catabolic traits. Researchers over the years has demonstrated the abilities of different group of microorganisms including fungi [10-12], bacteria [7,13,14,15], yeast [16] and algae [17-19] to degrade PAHs by pure cultures or when applied as consortium.

Among the microbes identified as contaminant degraders, the bacteria *Rhodococcus* and *Pseudomonas* have been frequently described in the catabolism of a range of substrates such as short-chain, long-chain, and halogenated hydrocarbons, and aromatic compounds, like PAHs, polychlorinated biphenyls and dibenzothiophenes [20-23]. Their well-established cellular resistance and metabolic ability for the degradation of all these compounds are related to their genomic properties, with an uncommon presence of multiple homologs of enzymes participating in major catabolic pathways and also a remarkable capacity for acquiring large linear plasmids which possibly conferred antibiotic resistance [20,24]. Their oxidative metabolism is capable of using several organic compounds as sole carbon and energy sources that fostered several industrial and bioremediation applications [20]. For the genus *Rhodococcus* the range of PAH serving as substrates includes fluorene, fluoranthene, phenanthrene, pyrene, chrysene, benz[a]anthracene, benzo[b]fluoranthene and benzo[a]pyrene [25-28]. Furthermore, metabolic pathway during biodegradation of PAHs by *Rhodococcus* spp has also been proposed from several previous reports [29,30].

The abilities of microbes to degrade both low molecular weight (LMW)-PAHs and HWM-PAHs has been extensively studied [26,29,31,32]. Bacterial degradation of PAH aerobically begins with the addition of both molecules of molecular oxygen to the aromatic ring by a ring hydroxylating dioxygenase (RHD), [33]. The RHD are multicomponent enzymes that consist of an electron transport chain, containing a ferredoxin and a reductase, and a terminal dioxygenase [34]. The latter is composed of two polypeptides, one of which (alpha subunit, RHD α) is the catalytic component of the enzyme system. The RHD α contain two conserved regions: the [Fe₂-S₂] Rieske center and the mononuclear iron-containing catalytic domain [35]. The RHD α serve as biomarkers of potential PAH-degraders, and the cognate genes can be PCR-amplified by using different primer sets [26,29,32,36].

In environments polluted by PAH, these compounds are invariably introduced in substances containing mixtures of PAH, such as coal tar and crude oil [37-40]. It's well-established that biodegradation of any given PAH can vary greatly when present as a part of a PAH mixture [39,41-

43], as other PAH may exert either positive or negative effects on biotransformation [37,43]. A variety of processes can operate simultaneously at cellular and intercellular (community) levels to contribute to variation of PAH biodegradation in PAH mixtures. It's possible that prolonged exposure to PAH mixtures can select for organisms adapted (e.g., enzyme specificities, regulatory circuits etc.) to catabolize PAH when present as mixtures. Thus, in efforts to identify microbes that effectively degrade PAH mixtures, employing PAH mixtures as enrichment substrates could be a strategy to acquire organisms better suited to function as bioremediation agents.

In this study, we examined the utility of using PAH mixtures, rather than single PAH, as enrichment substrates. We wished to determine if this approach would yield isolates that have superior PAH degradation abilities and retrieve isolates from diverse microbial taxa. We examined sediments of Lagos Lagoon, Nigeria, an important estuary in Africa that is impacted by PAH pollution [44].

Materials and Methods

Chemicals and reagents.

Analytical standards of various PAHs used in this study were obtained from Sigma Aldrich Corporation (St Louis, MO, USA). This includes fluorene (99% purity), pyrene (98% purity), benz[a]anthracene (99% purity) and fluoranthene (98% purity). HPLC grade organic solvents such as dichloromethane and acetone were purchased from Fisher Scientific Co. (Springfield, NJ, USA) while ethyl acetate was procured from Sigma Aldrich. Also purchased from Sigma Aldrich are analytical grades of the following compound; biphenyl, salicylic acid, phthalic acid, dibenzofuran, dibenzothiophene, methyl-naphthalene, acenaphthene, 2,4-dihydroxybenzoic acid, 2,5-dihydroxybenzoic acid, 3,4-dihydroxybenzoic acid, vanillic acid and 2,3-dihydroxybenzoic acid. Stock solutions of the hydrocarbon substrates were prepared by dissolution in acetone.

Sediment Sample Collection, Enrichment and Isolation of PAH-Degrading Bacteria

Sediment samples used for the enrichment technique were collected at three sampling stations (Apapa, Ofin and Eledu) from the Lagos Lagoon as described by Obi et al [44]. The continual enrichment technique adopted for the isolation of PAH-degrading bacteria has been described elsewhere [16]. Briefly, Sediment from each of the three sampling locations was mixed in equal parts to create a composite, and then 1 g of composite was suspended in

90 ml basal mineral salt medium (BMS) containing (g/l) 0.5 g (NH₄)₂SO₄, 1.0 g K₂HPO₄, 2.79g NaH₂PO₄, 0.2 MgSO₄·7H₂O, 0.5g NaCl, 50 mg Ca(NO₃)₂·4H₂O and 1mL of trace elements, pH 7.0±0.2. The medium was supplemented with the following PAHs mixture as sole sources of carbon and energy: pyrene, fluoranthene, benz[a]anthracene and fluorene at concentration of 25 mg l⁻¹ each making a total of 100 mg l⁻¹. The flasks were incubated at 30°C on an orbital shaker (Excella E10, New Brunswick Scientific Edison, New Jersey, USA) programmed at 150 rpm. After 14 days incubation, an aliquot of the enriched cultures was transferred into a fresh medium and handled as described above. This step was repeated 6 times to attain well-adapted PAH-degrading enriched bacterial consortia.

Bacterial strains in each consortium were isolated by spreading onto MSM-agar plates coated with a thin layer of different PAHs (benz[a]anthracene, pyrene, fluoranthene and fluorene) as described by Kiyohara et al. [45]. Bacterial colonies surrounded by a clear zone were scored positive, and subsequently purified and screened for utilization of PAHs.

Culture Maintenance

Pure cultures were grown overnight in Luria Bertani medium at 30°C, harvested by centrifugation (Eppendorf 5424/5424R Thermo Fisher Scientific, USA) at 6,000 rpm for 20 min. The pellets were resuspended in LB broth and subsequently distributed (1 ml) into 3 ml cryotubes already containing 1 ml glycerol (30% v/v). The mixture was thoroughly homogenized by shaking while avoiding foaming. The cryotubes were all stored at -20°C. For inocula development, isolates were routinely cultured on LB agar for purity check. Overnight cultures on LB broth were harvested by centrifugation, washed twice with MSM and subsequently resuspended in the same medium and adjusted to an OD (λ600 nm) of 1.

Hydrocarbon Degradation Profiling of Isolates

In addition to the PAHs of interest, isolates were tested for their abilities to grow on a range of other organic compounds (Table 1). Flasks containing 50 ml MSM were supplemented with 1% (v/v) petroleum feedstocks such as diesel, crude oil, pyridine and fresh engine oil. All other compounds were supplied at 100 mg l⁻¹. The flasks were inoculated with 200 μl of washed cell suspension and incubated on a rotary shaker at 30°C for 7 d in the dark.

PAH Biodegradation Studies

Pyrene, fluoranthene, benz[a]anthracene and

Table 1: Growth substrate range test.

+++ = Luxuriant growth (≥ OD600 = 1); ++ = Moderate growth (OD600 = 0.5-0.7), + = Slight growth (OD600 = 0.2-0.4); - = No growth.

		Isolate
Substrate	FF2	F4b
Pyrene	+++	+++
Phenanthrene	++	+
Fluorene	++	+++
Fluoranthene	+++	-
Benz[a]anthracene	+++	-
Methyl Naphthalene	+	-
2,4-Dihydroxybenzoic acid	+++	+++
2,5-Dihydroxybenzoic acid	+++	+++
3,5-Dihydroxybenzoic acid	+++	+++
Salicylic acid	++	++
Phthalic acid	++	+++
Vanillic acid	+++	+++
Acenaphthene	++	-
Dibenzofuran	+++	+
2,3-Dihydroxybenzoic acid	+	+++
3-Hydroxybenzoic acid	-	-
Pyridine	-	-
Biphenyl	+	++
Diesel	+++	+++
Crude oil	+++	+++

fluorene were added to sterile 250-ml conical flasks from concentrated stocks dissolved in acetone (100 ul of each). The acetone was dried under a sterile stream of nitrogen gas, and BMS (50 ml) was dispensed into the flask to obtain a final PAH concentration of 100 mg l⁻¹. For the degradation of PAH mixtures, the experiment was handled as described above except that 25 mg l⁻¹ of each the four PAH was added into a single 250 ml conical flask to obtain a final concentration of 100 mg l⁻¹. The flasks were inoculated with 200 μl of washed cell suspension and incubated at 30°C for 21 to 30 d on a rotary shaker at 180 rpm. Non-inoculated flasks served as negative controls for abiotic loss. Cell density (λ600 nm) and residual substrate concentration

were determined at each time point as biodegradation indices. All the treatments were established in triplicates.

Analytical Methods

Residual PAH concentration from both experimental and control flasks were analyzed by HPLC (Agilent, Santa Clara, CA) fitted with a programmable UV detector (UVD 170 U Dionex). Separations were performed on C-18 reversed-phased column (54 cm by 35 cm, i.d.; 4 μ m particle diam.) by isocratic elution with a mobile phase constituting 90% acetonitrile set at a flow rate of 1 ml min⁻¹. Absorbance of PAHs was monitored at 254 nm. The equipment was connected to a computer equipped with Chromeleon chromatography management software version 6.6. The software was used for quantification of PAHs throughout the experiment. PAH concentrations in the samples were determined against standard curves and the percentages of PAH remaining in the samples were calculated with the aid of deuterated PAH internal standard as a correction factor.

Genomic DNA Extraction

Genomic DNA of the PAH-degrading isolates was extracted using the UltraClean Microbial DNA Isolation Kit (MoBio Laboratories). The DNA was quantified using a fluorometer (Qubit[®] 3.0, ThermoFisher Scientific) and dsDNA quantitation done using the broad range protocol according to the manufacturer's instruction.

PCR Amplification and Sequencing of 16s Rrna Genes

PCR Amplification of the 16S rRNA coding region of the genomic DNA was performed using universal primers 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-GGTACCTTGTTACGACTT-3') (Akkermans et al., 1996). PCR was prepared in 50 μ l reaction mixtures containing 2.5 mM MgCl₂, 0.2 mM dNTPs, 50 pmol of each primer, 50 ng DNA template, 2.5 U Taq DNA polymerase (Promega) and PCR buffer (Promega, Madison WI). Amplification conditions used had an activation time of 3 min at 94°C, 30 cycles at 94°C for 1 min, 54°C for 1 min and 72°C for 1min, followed by 7 min of final primer extension at 72°C. Purified PCR products were cloned into the T-easy vector (Promega, Madison WI) and transformed into E. coli cells for propagation. Plasmids were extracted by using a plasmid miniPrep kit (Promega, Madison WI), and sequencing was done by the Sanger sequencing. Taxonomic identification was done by BLAST-N against the NCBI nr database. Multiple sequence alignments were created with MUSCLE and phylogenetic trees were constructed using MEGA 7 by the neighbor-joining algorithm and Kimura 2-parameter distance estimation method with bootstrap analysis for

1,000 replicates [46].

PCR amplification and sequencing of Pah-Rhd α genes.

Degenerate primers were used in PCR to amplify the PAH-RHD α in the isolates. For the Gram-positive bacteria these were Nid forward and Nid rev1 [36] while for Gram-negative bacteria, the primers were AC114F and 1096R [47]. PCR reactions were done in 50- μ l reaction mixtures containing 2.5 mM MgCl₂, 0.2 mM dNTPs, 500 nmol of each primer, 2.5 U Phusion polymerase (New England Biolabs) and PCR buffer (Promega, Madison WI). Amplification conditions were as follows: initial denaturation for 3 min at 94°C, 40 cycles at 94°C for 45 sec, 55°C for 45 sec and 72°C for 45 sec, followed by 7 min of final primer extension at 72°C. PCR reactions were validated by using genomic DNA extracted from Mycobacterium vanbaaleeni PYR1 and Delftia acidovorans Cs1-4 in which PAH metabolism and PAH degradation genes have been characterized. PCR products were resolved on 1.5% agarose gel and the band of expected size was excised, cleaned (Qiagen), cloned into TOPO 11 (ThermoFischer) and transformed into E. coli Top ten cells (ThermoFischer). Templates were prepared from colonies by amplification with the Templiphi system (GE Healthcare) and then submitted to the University of Wisconsin-Madison Biotechnology Center for sequencing. Vector and primer regions were trimmed from the sequences, which were then translated, and the resulting deduced amino acid sequence used for identification via BLASTP against the GenBank database. Multiple sequence alignments were created with MUSCLE and phylogenetic trees were constructed using MEGA 7 by the neighbor-joining algorithm and Kimura 2-parameter distance estimation method with bootstrap analysis for 1,000 replicates [46].

Nucleotide Sequence Deposition

The gene sequences determined in this study have been deposited in the GenBank database under accession numbers KT894768 (strain FF2) and KT894767 (strain F4b) for 16S rRNA genes and under accession numbers KT894771 (strain F4b) and KT894770 (strain FF2) for PAH-RHD α genes.

Results and Discussion

Isolation and Taxonomic Identification of Bacterial Isolates

The enrichment and isolation procedure produced a total of 12 bacterial isolates from growth on different PAHs (pyrene, fluoranthene, fluorene and benz[a]anthracene) and the mixtures of the four PAHs. Two strains (FF2 and F4b) exhibiting broad spectrum PAH utilization were selected for further studies. Strain FF2 showed a zone of clearance when

cultured on minimal salts agar sprayed with a thin film of each of the four test PAH added individually as well as in the mixture. However, while strain F4b created a zone of clearance when sprayed with fluorene, pyrene and the PAH mixture, no clearance was observed with fluoranthene and benz[a]anthracene. These findings indicate that clearance of PAH mixtures on spray plates can occur even if all of the PAH present in the mixture are not metabolized by the bacteria forming the colony. In this case, clearance may occur when surfactants released dissolve all PAH present.

The 16S rRNA gene of strain FF2 had 99% identity across the entire sequence (1476 bp) with those from *Rhodococcus* sp. strain 2G and *Rhodococcus pyridinivorans* strain SB3094. Construction of an evolutionary tree for strain FF2 showed that it clustered with three type strains *Rhodococcus* rhodochrous strain 372, *Rhodococcus biphenylivorans* strain TG9 and *R. pyridinivorans* strain PDB9, most closely with the latter (Figure 1). Hence, designation of this isolate as *R. pyridinivorans*. The description of the type strain *R. pyridinivorans* sPDB9, included growth on pyridine (the compound on which it was enriched, Yoon et al., [48] as a key chemotaxonomic characteristic. However, *R. pyridinivorans* strain FF2 did not utilize pyridine as a growth supporting substrate (see below), a discrepancy that likely reflects the fluidity of biodegradation genotypes relative to bacterial phylotypes.




Figure 1: Phylogenetic tree indicating the relationship of the 16S rRNA gene sequence from *R. pyridinivorans* strain FF2. Bootstrap values at given at the nodes. *Corynebacterium diphtheriae* NCTC11397 was used as the out group. The Genbank accession numbers for each sequence is given in parentheses.

For the strain F4b, construction of an evolutionary tree placed it within the *Pseudomonas aeruginosa* group (Figure 2). The species *P. aeruginosa* represents one of two primary lineages within the genus *Pseudomonas* (Figure




Figure 2: Phylogenetic tree indicating the placement of the 16S rRNA gene sequence from *P. aeruginosa* strain F4b within the genus *Pseudomonas*. Bootstrap values at given at the nodes. *Cellvibrio japonicus* Ueda107 was used as the out group. The Genbank accession numbers for each sequence is given in parentheses.

2). The other *Pseudomonas* lineage is the “*Pseudomonas fluorescens* complex”, which is constituted by several groups that collectively contains essentially all other *Pseudomonas* species except *P. aeruginosa* [49]. Thus, the clustering of the current isolate with *P. aeruginosa* type strains strongly supports the species designation for strain F4b.

Biodegradation of individual PAH

Strain FF2 grew with all four of the PAH tested when supplied individually (Table 1). Strain FF2 was able to utilize 82 % of the added benz[a]anthracene after 21 days of incubation, the rapid utilization of this compound is a characteristic that might make strain FF2 potentially useful in the bioremediation of PAH-contaminated sites, as removal of especially hazardous PAH like benz[a]anthracene is a key regulatory goal.

Strain F4b grew with pyrene and fluorene when supplied individually, but not with either benz[a]anthracene or fluoranthene (Table 1). The inability of *P. aeruginosa* strain F4b to utilize benz[a]anthracene was not unexpected because biodegradation of benz[a]anthracene by species of *Pseudomonas* are uncommon [50,51]. There are a few reports on biodegradation of pyrene by *Pseudomonas* species [52,53], but information on the biodegradation of other HMW PAH by pseudomonads is still limited.

Biodegradation of PAH in Mixtures

The abilities of strains FF2 and F4b to degrade mixtures of PAH are shown in (Figures 3 & 4), respectively. Strain FF2 utilized 70 %, 42 %, 35 % and 30 % of fluorene, pyrene, fluoranthene and benz[a]anthracene, respectively (Fig. 1). Strain F4b was able to degrade 10 % benz[a]anthracene, 50 % pyrene, 72 % fluorene and 30 % fluoranthene (Fig. 2)




Figure 3: Time course degradation of a mixture of pyrene, fluoranthene, fluorene and benz[a]anthracene degradation by *R. pyridinivorans* strain FF2




Figure 4: Time course degradation of a mixture of fluoranthene, pyrene, fluorene and benz [a]anthracene by *P. aeruginosa* strain F4b during biodegradation experiment.

after 21 days of incubation. In the presence of mixed PAH, *R. pyridinivorans* FF2 exhibited a specific growth rate and mean generation time of 0.072 h⁻¹ and 8.75 h, respectively (Table 2). *P. aeruginosa* strain F4b exhibited a specific growth rate of 0.0601 h⁻¹ and mean generation time of 7.64 h (Table 1).
Table 2: Growth kinetic parameters of strain FF2 and strain F4b.

T = Mean generation time, μ = Specific growth rate, NA = Not applicable

Substrates	Strain FF2		Strain F4b	
	T (h)	μ (hr ⁻¹)	T (h)	μ (hr ⁻¹)
Pyrene	6.993	0.09912	7.919	0.08753
Fluorene	9.008	0.07695	8.961	0.07735
Fluoranthene	10.74	0.06455	NA	NA
Benz[a]anthracene	14.46	0.04794	NA	NA
Mixed PAH	8.75	0.07213	7.64	0.07735

The greater efficacy of *R. pyridinivorans* strain FF2 in PAH biodegradation compared to that of *P. aeruginosa* strain F4b was consistent with prior reports. For example, Guo and coworkers [6] investigated the degradation of mixed PAHs by *Sphingomonas*, *Rhodococcus*, *Paracoccus* and *Mycobacterium* strains isolated from mangrove sediments. They observed that all of their isolates degraded completely a mixture of phenanthrene, fluoranthene and pyrene in 14 days. But, the Firmicute isolates (*Mycobacterium* and *Rhodococcus*) were more effective in degradation of mixed PAHs than were the Proteobacteria isolates (*Sphingomonas* and *Pseudomonas*). Mueller et al. [54] also concluded that biodegradation of PAH mixtures was more effective by Firmicutes than by Proteobacteria.

Growth of strain F4b in the presence of the four PAH resulted in degradation of 10% of benz[a]anthracene, which was likely cometabolism driven by the growth-supporting PAH, pyrene and fluorene (Figure 2). Juhasz et al. [55] reported cometabolic biodegradation of two HMW PAH, dibenz[a,h]anthracene and benzo[a]pyrene, by a *Burkholderia cepacia* strain utilizing phenanthrene for growth. Decreases of 41% and 52% in dibenz[a,h]anthracene and benzo(a)pyrene, respectively, were observed after 56 days of incubation. But, this organism could not grow on these compounds as a sole source of carbon and energy. Thus, degradation of dibenz[a,h]anthracene and benzo[a]pyrene by this *B. cepacia* strain was cometabolic, a case similar to the present finding of benz[a]anthracene degradation by *P. aeruginosa* strain F4b.

Substrate Utilization Specificity of Isolates

The abilities of the isolates to utilize a range of different hydrocarbons as carbon sources is summarized in (Table 1). The two isolates grew well on crude oil, engine oil, 2,5-dihydroxybenzoic acid, 2,4-dihydroxybenzoic acid, 3,5-dihydroxybenzoic acid and vanillic acid. Strain FF2 also grew well on benz[a]anthracene, fluoranthene and didenzofuran while it had moderate growth on fluorene, salicylic acid, phthalic acid and acenaphthene. Strain FF2 was not able to utilize 3-hydroxybenzoic acid or pyridine as sole carbon and energy sources. Strain F4b had good growth on fluorene, phthalic acid and 2,3-hydroxybenzoic acid and moderate growth on salicylic acid and biphenyl.

Overall, *R. pyridinivorans* strain FF2 exhibited a relatively broad spectrum of hydrocarbon utilization. Strong biodegradation capabilities of *R. pyridinivorans* strain FF2 mirror results reported by other investigators with other *Rhodococcus* isolates [20,38,56,57]. Pizzul et al. [58] reported that *Rhodococcus wratislaviensis*, degraded >

90 % of pyrene after 49 days while *Rhodococcus opacus* degraded 20 % of pyrene within a 10 day incubation period. The genus *Rhodococcus* exhibits a diverse range of metabolic activities, including the degradation of various aromatic and aliphatic hydrocarbons (short- and long-chain alkanes), monoaromatics (halogenated and nitro-substituted), heterocyclic compounds and polycyclic aromatic compounds [57].

P. aeruginosa strain F4b did not show growth when cultured in a minimal salts medium supplemented with methyl naphthalene, acenaphthene or pyridine, but grew well on fluorene, phenanthrene, phthalic acid, crude oil, diesel fuel, vanillic acid and other aromatic acids (Table 1). A similar finding was reported by Obayori et al. [53]. The luxuriant growth of *P. aeruginosa* strain F4b in the presence of crude oil and diesel fuel was important due to the fact that both contain a variety of hydrocarbons making it more likely to support microbial growth in actual contaminated environments [58]. Ilori and Amund [59] reported that *P. aeruginosa* isolated from a tropical soil polluted with crude petroleum was able to utilize crude oil and a host of other hydrocarbons.

For a microbe to be useful in hydrocarbon bioremediation, utilization of a wide range of hydrocarbons is a desirable characteristic [60] as these pollutants typically occur in the environment in complex mixtures. In this study, the two isolates showed a wide range of substrate utilization when grown in minimal salts medium (MSM) supplemented with different hydrocarbon substrates (Table 1).

Identification of PAH-RHDA genes.

For strain F4b the nahAC primers yielded a PCR product of expected size (ca. 500 bp), which had 100 % identity to nahAC-like genes of several *Pseudomonas* strains. Construction of a phylogenetic tree placed the PAH-RHD α of strain F4b within the nahAC group of the Gram-negative PAH-RHD α (Figure 5). For strain FF2 the nid primers also gave a product of the expected size (ca. 600 bp), which had 100 % identity to narAa of other *Rhodococcus* species/strains [61,62]. In the phylogenetic tree, the PAH-RHD α of strain FF2 was placed within narAa group of the Gram-positive PAH-RHD α (Figure 5).

Genes encoding PAH-RHD α are reliable biomarkers of PAH-degrading bacteria and their phylogeny provides a framework to distinguish genotypes based on PAH metabolism. The PAH-RHD α of Gram-negative and Gram-positive bacteria form two distinct lineages, each of which is comprised of several groups (Fig. 5). Within the Gram-




Figure 5: Phylogenetic tree of PAH-RHD α amplified from *R. pyridinivorans* strain FF2 and from *P. aeruginosa* strain F4b. The tree illustrates the main groups within the Gram-positive branch (*pdoA2/phdA*, *nidA3/fadA1*, *nidA/pdoA1*, *narAa*) and Gram-negative branch (*nahAC*, *nagAc*, *phnAC*, *ahdA1*). The Genbank accession numbers for each sequence is given in parentheses following each sequence. The Roman numerals and letters in parentheses following the groups are the clades corresponding to Meynet et al. (2015).

negative branch, the nahAC group is the largest and has been associated with PAH degraders spanning two classes, the gamma- and beta-proteobacteria (Cébron et al., 2008), and is invariably the genotype of PAH-degraders within the genus *Pseudomonas* [63-66]. Thus, identification of *P. aeruginosa* strain F4b as a nahAC-type PAH degrader is consistent with this framework. For strain FF2, the classification as narAa-type was also consistent with expectations as it is the genotype commonly associated with PAH degrading strains of *Rhodococcus* [63-66].

Conclusions

The use of multiple PAH during enrichment was effective for recovering bacteria with capabilities to grow on multiple PAH. The technique was successful in retrieving isolates from both the Proteobacteria and Firmicutes, and was thus capable of recovering isolates from a diverse range of microbial taxa. The present study extends our knowledge of PAH biodegradation by *P. aeruginosa* and is the first report of PAH biodegradation by *R. pyridinivorans*. The capability of the *R. pyridinivorans* isolate to effectively degrade a highly toxic PAH, benz[a]anthracene, has particular importance for use in bioremediation [67,68].

Acknowledgments

This study was supported by graduate fellowships from the University of Lagos and by an endowment from the O.N. Allen Professorship in Soil Microbiology to William J Hickey, University of Wisconsin-Madison, USA.

References

1. Cui M, Chen Y, Tian C, Zhang F, Yan C, et al. (2016) Chemical composition of PM_{2.5} from two tunnels with different vehicular fleet characteristics. *Sci. Total Environ* 550: 123-132.
2. Ravindra K, Sokhi R, Van Grieken R (2008) Atmospheric polycyclic aromatic hydrocarbons: source attribution, emission factors and regulation. *Atmos Environ* 42(13): 2895-2921.
3. Zhang Y N, Yang X L, Bian Y R, Gu C G, Liu Z T, et al. (2015) Aging law of PAHs in contaminated soil and their enrichment in earthworms characterized by chemical extraction techniques. *Huanjing Kexue/Environ. Sci* 36(12): 4582-4590.
4. Obi C C, Adebuseye S A, Ugoji E O, Ilori M O, Amund O O, et al. (2016) Microbial communities in sediments of Lagos Lagoon, Nigeria: Elucidation of community structure and potential impacts of contamination by municipal and industrial wastes. *Front Microbiol* 7, 1213.
5. Beriro D J, Cave M R., Wragg J, Thomas R, Wills G, et al (2016) A review of the current state of the art of physiologically-based tests for measuring human dermal in vitro bioavailability of polycyclic aromatic hydrocarbons (PAH) in soil. *J Hazard Mat* 305: 240- 259.
6. Guo C, Dang Z, Wong Y, Tam N Y F (2010) Biodegradation ability and dioxygenase genes of PAH-degrading *Sphingomonas* and *Mycobacterium* strains isolated from mangrove sediments. *Int Biodet Biodegrad* 64(6): 419-442.
7. Lyu Y, Zheng W, Zheng T, Tian Y (2014) Biodegradation of polycyclic aromatic hydrocarbons by *Novosphingobium pentaromativorans* US6-1. *Plosone* 9 (7): 1-8.
8. Haritash A K, Kaushik C P (2009) Biodegradation aspects of Polycyclic Aromatic Hydrocarbons (PAHs): A review. *J Hazard Mat* 169: 1-15.
9. Peng R H, Xiong A S, Xue Z, Fu X Y, Gao F, Zhao W, et al (2008) Microbial biodegradation of polyaromatic hydrocarbons. *FEMS Microbiol Rev* 32(6): 927-955.
10. Al-Hawash AB, Alkooranee JT, Zhang X, Ma F (2018) Fungal Degradation of Polycyclic Aromatic Hydrocarbons. *Int J Pure App Biosci* 6 (2), 8-24.
11. Moa J, Guan W (2016) Fungal degradation of polycyclic aromatic hydrocarbons (PAHs) by *Scopulariopsis brevicaulis* and its application in bioremediation of PAH-contaminated soil. *Acta Agriculturae Scandinavica, Section B, Soil Plant Sci* 66 (5): 1-7.
12. Mineki S, Suzuki K, Iwata K, Nakajima D, Goto S (2014) Degradation of polyaromatic hydrocarbons by fungi isolated from soil in Japan. *Polycycl. Aromat. Comp* 35(1): 120-128.
13. Darmawan R., Nakata H, Ohta H, Niidome T, Takikawa K, et al (2015) Isolation and Evaluation of PAH Degrading Bacteria. *J. Bioremed. Biodegrad* 6(3): 283.
14. Klankeo P, Nopcharoenkul W, Pinyakong O (2009) Two novel pyrene degrading *Diaphorobacter* sp. and *Pseudoxanthomonas* sp. isolated from soil. *J. Biosci Bioeng* 108(6): 488-495.
15. Kim Y H, Engesser K H, Cerniglia C E (2005) Numerical and genetic analysis of polycyclic aromatic hydrocarbon degrading *Mycobacteria*. *Microbiol. Ecol* 50(1): 110-119.
16. Hesham A, Khan S, Tao Y, Li D, Zhang Y, Yang M (2012) Biodegradation of high molecular weight PAHs using isolated yeast mixtures: application of meta-genomic methods for community structure analyses. *Environ Sci Pollut Res* 19(8): 3568–3578.
17. Takáčová A, Smolinská M, Ryba J, Mackuřák T, Jokrllová J, et al (2014) Biodegradation of Benzo[a]pyrene through the use of algae. *Cent. Eur J Chem* 12(11): 1133-1143.
18. El-Sheekh M M, Ghareib M M, EL-Souod G W.A (2011) Biodegradation of phenolic and polycyclic aromatic compounds by some algae and Cyanobacteria. *J. Bioremed Biodegrad* 2: 133.
19. Chan S M N, Luan T, Wong M H, Tam N F Y (2006) Removal and biodegradation of polycyclic aromatic hydrocarbons by *Selenastrum Capricornutum*. *Environ. Toxicol. Chem* 25(7): 1772–1779.
20. Rubashko G E, Kolomytseva M P, Golovleva L A (2006) Improvement of the Process of Fluorene Degradation by *Rhodococcus rhodochrous* strain 172. *Appl Biochem Microbiol* 42(4): 396–398.
21. de Carvalho CCR, Parreño-Marchante B, Neumann G, da Fonseca M M R, Heipieper H J (2005) Adaptation of *Rhodococcus erythropolis* DCL14 to growth on n-alkanes, alcohols and terpenes. *Appl. Microbiol. Biotechnol* 67(3): 383–388.
22. Abbey F W, Nicholas W H, Beshr F S, Kathrine C H, Lee D (2011) Remediation of PAH-Contaminated Soils and Sediments: A Literature Review. pp. 1-102.
23. Abalos A M, Vinãas J, Sabate´ M A, Solanas M A (2004) Enhanced biodegradation of Casablanca crude oil by a microbial consortium in presence of a rhamnolipid produced by *Pseudomonas aeruginosa* AT10. *Biodegra* 15(4): 249-260.
24. Van der Geize R, Dijkhuizen L (2004) Harnessing the catabolic diversity of rhodococci for environmental and biotechnological applications. *Curr Opin Biotechnol* 7(3): 255-261.
25. Yu C., Yao J, Jin J (2014) Characteristics and metabolic pathways of fluorene (FLU) degradation by strain *Rhodococcus* sp. USTB-C isolated from crude oil. *J Chem Pharm Res* 6 (3): 560-565.
26. Mishra S, Singh S N (2014) Biodegradation of benzo(a)pyrene mediated by catabolic enzymes of bacteria *Int J Environ Sci Technol* 11: 1571–1580
27. Subashchandrabose SR, Venkateswarlu K, Naidu R, Megharaj M (2019) Biodegradation of high-molecular weight PAHs by *Rhodococcus wratislaviensis* strain 9: Overexpression of amidohydrolase induced by pyrene and BaP. *Sci. Total Environ* 651 (Pt 1): 813-821.
28. Peng T, Luo A, Kan J, Liang L, Huang T, Hu Z J (2018) Identification of A ring-hydroxylating dioxygenases capable of anthracene and benz[a]anthracene oxidation from *Rhodococcus* sp. P14. *J Mol Microbiol Biotechnol* 28(4): 183-189.
29. Song X., Xu Y., Li G, Zhang Y, Haung T, Hu Z (2011) Isolation, characterization of *Rhodococcus* sp. P14 capable of degrading high-molecular-weight polycyclic aromatic hydrocarbons and aliphatic hydrocarbons. *Mar Pollut Bull* 62 (10): 2122-2128.
30. Dean-Ross D, Moody JD, Freeman JP, Doerge D R, Cerniglia CE (2001) Metabolism of anthracene by a *Rhodococcus* species. *FEMS Microbiol. Lett* 204: 205-211.
31. Zhou HW, Gu CL, Wong YS, Tam NFY (2006) Genetic diversity of dioxygenase genes in polycyclic aromatic hydrocarbon-degrading bacteria isolated from mangrove sediments. *FEMS Microbiol. Lett* 262: 148–157.

Citation: Chioma C Obi, Sunday A Adebuseye, Olukayode O Amund, Esther E Ugoji and William J Hickey. Biodegradation of Polycyclic Aromatic Hydrocarbon Mixtures by *Rhodococcus Pyridinivorans* FF2 and *Pseudomonas Aeruginosa* F4b Isolated from Sediments of Lagos Lagoon, Nigeria. *Op Acc J Bio Sci & Res* 2(5)-2020.

DOI: 10.46718/JBGSR.2020.02.000058

32. Demaneche S, Meyer C, Micoud J, Louwagie M, Willison J C, et al. (2004) Identification and functional analysis of two aromatic-ring-hydroxylating dioxygenases from a *Sphingomonas* strain that degrades various polycyclic aromatic hydrocarbons. *Appl. Environ. Microbiol* 70(11): 6714–6725.
33. Chadhain SM, Norman RS, Pesce KV, Kukor JJ, Zylstra GJ (2006) Microbial dioxygenase gene population shifts during polycyclic aromatic hydrocarbon biodegradation. *Appl. Environ. Microbiol* 72: 4078-4087.
34. Cerniglia C (1992) Biodegradation of polycyclic aromatic hydrocarbons. *Biodegrad* 3: 351-368.
35. Baldwin B R, Nakatsu C H, Nies L (2003) Detection and enumeration of aromatic oxygenase genes by multiplex and real-time PCR. *Appl. Environ. Microbiol* 69(6): 3350-3358.
36. Bordenave S, Goni-urriza M, Vilette C, Blanchard S, Caumette Duran R (2008) Diversity of ring-hydroxylating dioxygenases in pristine and oil contaminated microbial mats at genomic and transcriptomic levels. *Environ. Microbiol* 10: 3201-3211.
37. Hennessee CT, Li QX (2016) Effects of polycyclic aromatic hydrocarbon mixtures on degradation, gene expression, and metabolite production in four *Mycobacterium* species. *Appl. Environ. Microbiol* 82: 3357-3369.
38. Dean-Ross D Moody J, Cerniglia CE (2002) Utilization of mixtures of polycyclic aromatic hydrocarbons by bacteria isolated from contaminated sediment. *FEMS Microbiol. Ecol* 41(1): 1-7.
39. Brauner JS, Widdowson M A, Novak JT, Love NG (2010) Biodegradation of a PAH Mixture by Native Subsurface Microbiota. *Bioremed* 6(1): 9-24.
40. McNally LD, Mihelcic JR, Lueking DR (1999) Biodegradation of mixtures of polycyclic aromatic hydrocarbons under aerobic and nitrate-reducing conditions. *Chemosphere* 38(6): 1313-1321.
41. Lotfabad SK, Gray MR (2002) Kinetics of biodegradation of mixtures of polycyclic aromatic hydrocarbons. *Appl. Microbiol Biotechnol* 60(3): 361-366.
42. Desai AM, Christidus PD, McDonald TJ (2008) Biodegradation kinetics of select polycyclic aromatic hydrocarbon (PAH) mixtures by *Sphingomonas paucimobilis* EPA505. *Biodegrad* 19(4): 223-233.
43. Christidus PD, Autenrieth RL (2007) Kinetics of biodegradation of binary and ternary mixtures of PAHs. *Biotechnol Bioeng* 97 (4): 788-800.
44. Obi CC, Adebuseye SA, Amund OO, Ugoji EO, Ilori MO, et al. (2017) Structural dynamics of microbial communities in polycyclic aromatic hydrocarbon-contaminated tropical estuarine sediments undergoing simulated aerobic biotreatment. *Appl. Microbiol. Biotechnol* 101: 4299-4314.
45. Kiyohara H, Nagao K, Yana K (1982) Rapid screen for bacteria degrading water insoluble, solid hydrocarbons on agar plates. *Appl. Environ. Microbiol* 43: 454-457.
46. Kumar S, Stecher G, Tamura K. (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol. Biol. Evolut* 33(7): 1870-1874.
47. Wilson MS, Herrick JB, Jeon CO, Hinman DE, Madsen EL (2003) Horizontal transfer of *phnAc* dioxygenase genes within one of two phenotypically and genotypically distinctive naphthalene-degrading guilds from adjacent soil environments. *Appl Environ Microbiol* 69(4): 2172-2181.
48. Yoon JH, Kang SS, Cho YG, Lee ST, Kho YH, et al. (2000) *Rhodococcus pyridinivorans* sp. nov., a pyridine-degrading bacterium. *Int J Syst Evol Microbiol* 50(6): 2173-2180.
49. Garrido-Sanz D, Meier-Kolthoff JP, Göker M, Martín, M, Rivilla R, et al. (2016) Genomic and Genetic Diversity within the *Pseudomonas fluorescens* Complex. *PLoS ONE* 11 (2): 1371.
50. Kanaly RA, Harayama S (2010) Advances in the field of high-molecularweight polycyclic aromatic hydrocarbon biodegradation by bacteria. *Microb. Biotechnol* 3(2): 132-164.
51. Seo J S, Keum YS, Li QX (2009) Bacterial degradation of aromatic compounds. *Int. J. Environ. Res. Pub. Health* 6(1): 278-230.
52. Kazunga C, Aitken MD (2000) Products from the incomplete metabolism of pyrene by polycyclic aromatic hydrocarbon-degrading bacteria. *Appl. Environ. Microbiol* 66(5): 1917-1922.
53. Obayori OS, Ilori MO, Adebuseye SA, Amund OO, Oyetibo GO (2008) Microbial population changes in tropical agricultural soil experimentally contaminated with crude petroleum. *Afr J Biotechnol* 7: 4512-4520.
54. Mueller JC, Devereux R, Santavy DL, Lantz SE, Willis SG, et al. (1997) Phylogenetic and physiological comparisons of PAH-degrading bacteria from geographically diverse soils. *Antonie Van Leeuwenhoek* 71: 329-343.
55. Juhasz A L, Britz M L, Stanley V (1997) Degradation of fluoranthene, pyrene, benz[a]anthracene and dibenz[a,h]anthracene by *Burkholderia cepacia*. *J Appl Microbiol* 83: 189-198.
56. Pizzul J, Castillo MP, Stenström J (2007) Initial microbial degradation of polycyclic aromatic hydrocarbons. *Biodegrad* 59: 111-118.
57. Guo CL, Zhou HW, Wong YS, Tam NYF (2005) Isolation of PAH-degrading bacteria from mangrove sediments and their biodegradation potential. *Mar. Pollut. Bull* 51(8-12): 1054-1061.
58. Adebuseye S A, Ilori M O, Teniola O D, Olatope S O (2007) Microbial degradation of petroleum in a polluted tropical stream. *World J. Microbiol. Biotechnol* 23: 1149-1159.
59. Ilori MON, Amund D I (2000) Degradation of anthracene by bacteria isolated from oil polluted tropical soils. *Zeitschrift für Naturforschung* 55(11-12): 890.
60. Brooijmans RJW, Pastink MI, Siezen RJ (2009) Hydrocarbon-degrading bacteria: the oil-spill clean-up crew. *Microb. Biotechnol.* 2 (6L): 587–594.
61. Treadway SL, Yanagimachi KS, Lankeau E, Lessard PA, Stephanopoulos G, et al. (1999) Isolation and characterization of indene bioconversion genes from *Rhodococcus* strain I24. *Appl Microbiol Biotechnol* 51(6): 786-793.
62. Anan Ina LN, Yastrebova OV, Demakov VA, Plotnikova EG (2011) Naphthalene-degrading bacteria of the genus *Rhodococcus* from the Verkhnekamsk salt mining region of Russia. *Antonie Van Leeuwenhoek* 100 (2): 309-316.
63. Moser R, Stahl U (2001) Insights into the genetic diversity of initial dioxygenases from PAH-degrading bacteria. *Appl. Microbiol. Biotechnol* 55(5): 609-618.
64. Habe H, Omori T (2003) Genetics of polycyclic aromatic

Citation: Chioma C Obi, Sunday A Adebuseye, Olukayode O Amund, Esther E Ugoji and William J Hickey. Biodegradation of Polycyclic Aromatic Hydrocarbon Mixtures by *Rhodococcus Pyridinivorans* FF2 and *Pseudomonas Aeruginosa* F4b Isolated from Sediments of Lagos Lagoon, Nigeria. *Op Acc J Bio Sci & Res* 2(5)-2020.

DOI: 10.46718/JBGSR.2020.02.000058

-
- hydrocarbon metabolism in diverse aerobic bacteria. *Biosci. biotechnol biochem* 67 (2): 225-243.
65. Cébron A, Norini MP, Beguiristain T, Leyval C (2008) Real-time PCR quantification of PAH-ring hydroxylating dioxygenase (PAH-RHD α) genes from Gram positive and Gram negative bacteria in soil and sediment samples. *J Microbial Methods* 73(2): 148-159
66. Meynet P, Head IM, Werner D, Davenport RJ (2015) Re-evaluation of dioxygenase gene phylogeny for the development and validation of a quantitative assay for environmental aromatic hydrocarbon degraders. *FEMS Microb Ecol* 91(6): 1-11.
67. Kim SJ, Kweon O, Jones RC, Freeman JP, Edmondson RD (2007) Complete and integrated pyrene degradation pathway in *Mycobacterium vanbaalenii* PYR-1 based on systems biology. *J Bacteriol* 189(2): 464-472.
68. Kulakov LA, Allen CCR (2005) Biodegradation and *Rhodococcus* – masters of catabolic versatility. *Curr Opin Biotechnol* 16(3): 282–290.

*Corresponding author: : William J Hickey, Email: Amany_Alzokaky@yahoo.com

Next Submission with BGSR follows:

- Rapid Peer Review
- Reprints for Original Copy
- E-Prints Availability
- Below URL for auxiliary Submission Link: <https://biogenericpublishers.com/submit-manuscript/>