



Article Type: Review Article

Received: 14/08/2020

Published: 20/08/2020

DOI: 10.46718/JBGSR.2020.04.000094

Covid-19 Re-Infection among Recovered Patients and its Implication for Lung Health: A Review

Obande-Ogbuinya Nkiru Edith¹, Aleke Christian Okechukwu^{2*}, Omaka-Omari Lois Nnenna², Afoke Eunice Nwafor² and Nwankwo Onyechi³

¹Department of Science Education, Faculty of Education, Alex Ekwueme Federal University, Ndufu Alike, Ebonyi State Nigeria.

²Department of Human Kinetics and Health Education, Ebonyi State University, PMB 053 Abakaliki, Ebonyi State, Nigeria

³African Institutes for Health Policy and Health Systems, Ebonyi State University, Nigeria

*Corresponding author: Aleke Christian Okechukwu, Department of Human Kinetics and Health Education, Faculty of Education, Ebonyi State University, Abakaliki, Nigeria

Abstract

COVID-19 infection has continued to pose as a very serious health threats to mankind globally despite all efforts geared toward curbing its spread. More worrisome recently is the report from different parts of the world on re-infection of those treated and recovered with COVID -19 patients thus making containment of the virus even more difficult. Of more worrisome is the fact that the lung, a vital human organ is the major site being attacked by the virus even on re-infection cases. If quick action is not taken early enough, it may lead to outright death of the patient. A lung infection, (Pneumonia) caused by COVID-19 has been discovered to be having a stunning effect on hospital systems and killing COVID virus patients silently and it occurs even as the patient is asymptomatic. This paper examines the reasons for re-infection, Lacuna in the reviewed literature vis-à-vis PCR test results, effect of re-infection on the lungs and implication for patients' lung health. The papers summarized and concluded that it's a fact that re-infection occurs among patients accompanied with mild or severe symptoms having far reaching implications for the patient's lung health. The paper recommends amongst others that government at all levels should collaborate with WHO, CDC and health policy makers legally mandating, that every recovered patient should stay additional 2weeks in the hospital for early detection of re-infection in order to avert any invasion and damage to the lungs thus ensuring lung health. Also proper health education should be availed to the recovered patients to avoid any exposures or habits (different from the index disease) such as smoking that can pose dangers to the already fatigued lungs.

Keywords: COVID-19 Pneumonia; Coronavirus re-infection; recovered patient; Lung Health

Introduction

The novel coronavirus disease 2019 (COVID-19) pandemic outbreak, which started in the Hubei province of China in December 2019, has become a major public health challenge for not only China but also countries around the world. The COVID-19 pandemic is rapidly evolving and has since spread to all continents in the world affecting over 210 countries. Since 31 December 2019 and as of 01 May 2020, 3214256 cases of COVID-19 have been reported, including 232570 deaths and the number of reported cases is rapidly increasing worldwide [1].

COVID-19 infection has continued to pose as a very

serious health threats to mankind globally despite all efforts geared toward curbing its spread and consequential high mortality. More worrisome recently is the report from different parts of the world on re-infection of recovered patients (those who were infected with COVID -19 and got treated) thus making containment of the virus even more difficult [2]. Of more worrisome is the fact that the lung, a vital human organ is the major site being attacked by the virus even on re-infection cases. If quick action is not taken early enough, it may lead to outright death of the patient [2]. The pneumonia caused by the virus, COVID pneumonia was found to be the major cause of deaths in COVID-19 patients. It causes much oxygen deprivation (hypoxia) and

difficult to detect in nature [3]. The COVID pneumonia (lung infection) does not present early even when the patient has been infected and yet to complain of breathing difficulty. All these call for worry globally, continentally and nationally about cases of re-infection.

In utter dismay on cases of re-infection (where patients assumed to have been treated of COVID-19 get hit once again from the virus), Studies have reported that over 163 recovered COVID-19 patients in Korea who on recovery tested positive once again as revealed by Korean Centre for disease control (KCDC) [4]. Studies also observed patients who on seemingly recovery from COVID-19 tested positive to the virus, one hundred and eleven (111) of such cases were cited by South Korean Officials [5]. Additional information on recovered covid-19 patients in Southern Chinese city of Shenzhen revealed that 38 out of 262 (15%) of them tested and confirmed positive via Polymerase chain reaction (PCR). It is currently regarded as the gold standard for COVID-19 testing. The 38 patients were mostly young people below the age of 14 showed mild symptoms during their period of infection. In Wuhan China where the COVID-19 was first reported, a report was made of four medical workers who had 3 consecutive positive PCR tests after recovery. Just as the study in Shenzhen, the patients were asymptomatic and their family members not infected [5]. A 40-year old COVID-19 patient who had the RT-PCR examination for SARS-CoV-2 performed on her in day 10, 11 and 14 of admission tested negative to all the 3 tests and was discharged although asked to be on quarantine at home. On day 24, the patient was subjected to another test for SARS-CoV-2 having presented pain in the chest and production of cough and sputum. Surprisingly, the test turned out positive hence resulting to the patient being readmitted in Taiwan hospital hence the result showed absorption of lung disease [6].

Studies have reported cases in Japan of a woman who tested positive of COVID-19 the second time followed by China reports of numerous Chinese patients testing positive after discharge from the isolation centers. On a good note according to the authors, such patients were not found to be infectious. The authors posited that the convalescing COVID-19 patients may not have built enough antibodies or immunity to SARS-CoV-2 and thus become infected again. It could also be that the virus is biphasic like anthrax which entails the virus lying dormant before developing new symptoms. In China, some of the first cases of re-infection were attributed to discrepancies in testing. For instance, some hospitals tested samples from noses and throats before patients were discharged only to discover the virus in the lower respiratory tracts of the patients on re-infection. This means that patients may remain carriers

despite meeting discharged criteria [2].

Allen Cheng, a professor of infectious disease proclaimed that the details of the Japan cases put to rest the argument of whether the recovered patients tested positive to COVID-19 as a result of them being persistently positive on disappearance of symptoms or that the patients have been re-infected. It was conclusively put that the patients had been re-infected [7]. One good sign according to Song Tie on China disease control remains that none of the patients appeared to have infected other people after discharge. In other words, it means that the re-infected patient may not be contagious. It could also mean that specific antibodies developed by convalescing patients make them to be immuned to the virus which infected them. However, the infectious disease specialist observed that the second re-infection is mostly non severe [8]. A new study in China revealed that recovered patients usually have very low or even insignificant levels of antibodies to fight the virus. The study showed that a third of recovered patients in China had low levels of antibodies with only ten of the patients having no detectable antibody to neutralize the virus. This raises a lot of concern because when the recovered patients fail to develop high level immunity to ward off re-infection, it could adversely affect herd immunity needed to stop wider spread. This implies difficulty for development of a vaccine. When the real virus could not trigger an antibody response, of what role and use would a weakened version normally used in making vaccine be? [9]. The WHO nevertheless, is investigating more reports of re-infection among recovered patients [10]. However, the (Table 1) below shows a schematic framework on the effect of COVID-19 on the lungs.

Table 1: Symptoms of COVID-19.

Symptoms	COVID-19	Influenza (Flu)	Cold
Dry Cough	+++	+++	+
Fever	+++	+++	-
Stuffy nose	-	++	+++
Sore Throat	++	++	+++
Shortness of Breath	++	--	--
Headache	++	+++	-
Body Aches	++	+++	+++
Sneezing	--	--	+++
Exhaustion	++	+++	++
Diarrhea	-	++	--
+++Frequent ++Sometimes +Occasionally -Rare --Not Observed			
Source: WHO, CDC in Freund 2020			

Citation: Obande-Ogbuinya Nkiru Edith, Aleke Christian Okechukwu*, Omaka-Omari Lois Nnenna, Afoke Eunice Nwafor and Nwankwo Onyechi . Covid-19 Re-Infection among Recovered Patients and its Implication for Lung Health: A Review. Op Acc J Bio Sci & Res 4(2)-2020.

Reasons for Covid-19 Re-Infection Among Recovered Patients

Numerous reasons have been availed for the positive test results of recovered patients. It could be that virus load gets detected around 10 weeks of onset of covid-19 about the time some of the patients are discharged. It could also be that SARS-CoV2 sheds from its victims for a prolonged time. Also, since the viral load in a sample seems to be fluctuating during disease course, there may be high tendency for a negative result of covid-19 test even after treatment [4]. Another reason as reviewed could be that the convalescing COVID-19 patients may not have built enough antibodies or built up immunity to SARS-CoV-2 and thus become infected again. It could also be that the virus is biphasic like anthrax which entails the virus lying dormant before developing new symptoms [6]. Be it as it may, it is still unclear whether the second positive result of a treated COVID-19 patient can be linked to transmission of the infection to other persons. However, given to the disastrous outcome of imminent disease transmission, longer observation period ought to be taken into consideration for Covid-19 patients [6].

On the other hand, the presence of severe acute respiratory syndrome coronavirus 2(SARS-CoV-2) viral RNA in a respiratory specimen may not be directly taken as a possibility or probability for disease infection or transmission. In the words of the authors, it is a known fact that even after infection with some of the past viral diseases like Ebola virus, Middle East respiratory syndrome, Coronavirus, Zika virus, Influenza virus and Measles virus, that the viral RNA can still be seen long after the disappearance of the virus. For instance, the measles virus can still be seen 6-8 weeks even after clearance of the infectious virus. During the period, the immune system usually tries to neutralize viruses by envelop lysing or by bringing together (aggregating) the particles of the virus. The efforts are always in the bid to avoid or prevent re-infection. However, the RNA does not become eliminated. It rather gets degraded gradually and slowly over a period of time. Be that as it may, making use of sensitive PCR methods can offer value in diagnosing cases but caution should be observed as the PCR does not differentiate infectious virus from non infectious nucleic acid [11]. The study warned against the use of findings of their study in making conclusions about prolonged shedding of virus or even using their findings to make or amend policies on isolation hence more research or data on viral load is needed.

From the reviews, it can easily be summarized that it is a reality on ground that recovered patients can be re-infected. Other people may not be infected by a recovered re-infected patient. On re-infection, the symptoms may

appear mild or less severe than the first infection period. Antibodies are not in sufficient levels (detectable levels) or may even be unavailable at recovery. The re-infections as reviewed happen between 10 days and 2 weeks on recovery.

Lacuna in the Review Vis-À-Vis Pcr Test Results

The finding of the studies that the PCR does not differentiate infectious virus RNA from non infectious Virus RNA invariably contradicts the popular perception that re-infected patients cannot infect other people [11]. The infectious Virus RNA as the name depicts may infect others. This earnestly calls for a more in-depth study that will determine whether the infectious RNA can actually be contagious and capable of infecting others. This will avert further transmission of the virus from those already assumed to have fully recovered.

Covid-19 Re-Infection And Patient's Lung Health

The Lung is a vital organ of the body playing great roles in respiration, circulation, excretion and speech production in humans [12]. Any cause depriving the lungs from performing these functions has affected lung health. Lung health in the context of this paper means the lung being in a complete stable and disease free. Any damage to the lungs as a result of infection or re-infection has affected virtually every biological functioning of the body that may lead to eventual death. It is a known fact that COVID-19 leaves patient with severe lung problems. When the patient is re-infected, how is the secondary effect on lungs? Does it imply further damage to the lungs since it also comes with symptoms although mild in nature as pointed out from the reviewed studies? Answering these questions will be of significance to WHO, various countries' Centers for disease control, medical personnel and health policy makers. It will help them to develop and establish the quickest means and channel of treating the new re-infected cases so as to avert further fatal lung damage or death arising from complications experienced by already assumed recovered patients. The most convenient duration at most for a patient to be discharged can be unveiled when the generated questions are answered. It will definitely bring to a halt, the current norm of some recovered patients going home only to return few days after with COVID-19 re-infection symptoms. It will also bring to the barest minimum lung incapacity and irredeemable damage caused by re-infection.

Effects of Covid-19 on the Lungs of Re-Infected Patients: Implication for Lung Health

When the lungs become filled with inflammatory material during COVID-19 infection, they can no longer get sufficient oxygen to the blood stream thus reducing the body's ability to take oxygen around the body and getting

rid of carbon dioxide. It is the most cause of death due to severe pneumonia. Also when the virus reaches the lungs, the mucous membranes are inflamed leading to damage on the alveoli or lung sacs. They in turn try to work harder in order to carry out their oxygen supply function to the rest of the body parts and also removing carbon dioxide from the blood as to be exhaled. The swelling of liver lymph nodes caused by the virus can impair oxygen flow. It can also cause the lung areas to be filled with pus apart from fluid and dead cells hence the occurrence of pneumonia which is infection of the lung. In worst cases known as acute respiratory distress syndrome, the lungs may be so filled with much fluid that no amount of life support may be of help and the patient dies [13,14].

From the review, it implies that the virus reaching the lungs at any point in time does not say well for the lung's health whether first time infection or on re-infection. A lung infection mild or severe impedes normal breathing pattern and affects almost all vital body functioning waste management inclusive [15]. In either mild or severe lung infection, a prompt medical aid is needed to avert imminent mortality. The review has also pointed out that during re-infection of the lungs that may have been caused by re-infection with COVID-19, other parts of the body may first be infected before the lungs implying urgent medical attention on those parts before the Lungs can be reached. A lung infection with corona virus may also lead to infection of other cells such as the gastro intestinal system which explains the symptoms of diarrhea or indigestion in some patients. Studies have observed that at a re-infected COVID-19 patient may end up with reduced lung capacity and function following bouts of the disease [16-19]. It was discovered by a Hongkong hospital Authority that 2 or 3 recovering patients had lost 20-30% of lung function.

Conclusion

Based on the reviews, there have been cases of re-infection among recovered patients. The symptoms are the same as the first instance although less severe. Regardless of the mild symptoms on re-infection, the lungs may be invaded once more and sometimes left incapacitated or completely damaged. Lung health being a sinequanon for human existence, prompt medical attention ought to be quickly availed when patients present re-infection by COVID-19. Based on this the paper recommended that:

a. WHO in collaboration with CDC of different nations and health policy makers should make quick health policies backed by law, mandating every recovered patient from COVID-19 to be in hospital custody for at least two weeks upon discharge. This will also make for easy and early

detection of re-infection and afford the patient access to ventilators and other equipment for enhancement of lung health.

b. The Government at all levels should organize periodic free Lung function tests for recovered Corona virus patients for early detection and treatment of any ailing lung.

c. Health Sectors should collaborate with Manufacturers to produce and make free distribution of small and more portable ventilators for every COVID-19 Patient whether recovered or still infected for proper management of lung health.

d. Proper health education should be availed to the recovered patients to avoid any exposures or habits (different from the index disease) such as smoking that can pose dangers to the already fatigued lungs

References

1. (2020) European Centre for Disease Prevention and Control. An agency of the European Union. Situation update worldwide
2. Stanway D, Kelland K (2020) Explainer: Coronavirus reappears in discharged patients, raising questions in containment fight. *Coronavirus Bulletin*.
3. Levithan R (2020) The infection that's silently killing Corona virus Patients.
4. Hancocks P, Seo Y, Hollingsworth J (2020) Recovered coronavirus patients are testing positive again. Can you get reinfected? Cable News Network Turner Broadcasting System, Inc.
5. Leung H (2020) Can you be re-infected after recovering from Corona virus? Here's what we know about covid-19 immunity.
6. Zhou M, Li Q, Cao L, Liu Y, Zha Y et al. (2020) Re-emergence of SARS-CoV2 in a discharged COVID-19 case. *Journal of Microbiology, Immunology and Infection* 53(3): 501-502.
7. Cheng A (2020) Concerns as Covid-19 reappears in discharged patients worldwide *New Straits Times*, New Straits Times Press (M) Bhd. A part of Media Prima Group 2020
8. Tie S (2020) Discharged patients test positive again *The Star Online* 2020
9. Nisar ul H (2020) Patients who recover are getting infected with the virus again. *Asian web news on covid-19*.
10. Nebehay S (2020) WHO is investigating reports of recovered COVID-19 patients testing positive again.
11. Atkinson B, Petersen E (2020) SARS-CoV-2 shedding and infectivity. *The Lancet*. 2020.
12. Sampson S (2020) What do the lungs do? 2018.
13. Readfern G (2020) What happens to people's lungs when they get corona virus?, *The Guardian* 2020.
14. Belluk P (2020) What does the corona virus do to the body?
15. Bradford A (2020) Lungs : facts, function and diseases. 2018.
16. Healy M (2020) Coronavirus may cause lasting damage throughout the body, doctors fear.
17. McFall-Johnsen M (2020) More than 370, 000 people have recovered from COVID-19. Here's what we know about the

-
- Survivors.
18. Freund A (2020) COVID-19: recovered patients have partially reduced lung function. Doc's association Kashmir urges immune testing for recovered patients Asian Web News.
19. Pawlowski A (2020) What are the big-term health consequences of COVID-19? 2020.

*Corresponding author: : Aleke Christian Okechukwu,. E-mail: christian.aleke@yahoo.com; Christian.aleke2020@gmail.com

Next Submission with BGSR follows:

- Rapid Peer Review
- Reprints for Original Copy
- E-Prints Availability
- Below URL for auxiliary Submission Link: <https://biogenericpublishers.com/submit-manuscript/>