



Article Type: Research Article

Received: 19/03/2021

Published: 25/03/2021

DOI: 10.46718/JBGSR.2021.08.000190

The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on *Plasmodium berghei* in Swiss Albino Mice

Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰

¹Department of Zoology, Faculty of Science, Federal University of Lafia, Lafia, Nasarawa State, Nigeria.

²Department of Animal Biology (Parasitology and Tropical Diseases Research Unit), Federal University of Technology, Minna, Niger State, Nigeria

³Department of Toxicology and Pharmacology, National Institute of Pharmaceutical Research and Development, Federal Capital Territory, Abuja, Nigeria

⁴National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention, Shanghai, China

⁵Department of Chemistry, Faculty of Science, Federal University of Lafia, Lafia, Nasarawa State, Nigeria

⁶Department of Biological Science, Faculty of Science, Gombe State University, Gombe, Gombe State, Nigeria

⁷Department of Chemistry, School of Physical Science, Federal University of Technology Owerri, Imo State, Nigeria

⁸Faculty of Natural and Applied Sciences, Godfrey Okoye University, Enugu, Enugu State, Nigeria

⁹Department of Zoology, Faculty of Natural Sciences, University of Jos, Jos, Plateau State, Nigeria

¹⁰Department of Microbiology, Faculty of Natural Science, Plateau State University Bokoos.

*Corresponding author: Ayim JO, Department of Zoology, Faculty of Science, Federal University of Lafia, Lafia, Nasarawa State, Nigeria. Email: joeayim2017@gmail.com

Abstract

Malaria remains one of the most prevalent and ravaging infection across Africa especially in Nigeria. Malaria parasite success or proliferation in its host have been linked to the ability of the parasite to show resistance to diverse anti-malarial drugs, thereby leading to the utilization of natural products from animals and plants as a potential strategy for identifying novel antimalarial drugs. Thus, this study aimed at screening toad venom for its antiplasmodial potency in swiss albino mice. Toads were collected from July to August 2019. The acute oral toxicity and biochemical characterization of the toad venom was determined. The experimental mice was administered varying curative doses (130, 173 and 217 mg/kg) of the toad venom crude extract and standard drug once daily for 3 days. *in-vivo* antiplasmodial activity was evaluated by a curative test, after infecting the mice model with *Plasmodium berghei*. From the results obtained, parasitemia was highest in the control group while the group treated with Artemether-lumerfantrin (ACT) had the least parasitemia. Thus, the mean parasitemia in relation to the different treatments varied significantly ($P < 0.05$). Nevertheless, the mean parasitemia in albino mice in relation to the highest dosage of toad venom versus ACT treatments showed no significant difference ($P > 0.05$). The mean change in haemoglobin (Hb) and as well as pack cell volume (PCV) levels in albino mice after four days in relation to treatments with toad venom and ACT showed a significant difference ($P < 0.05$) whereas the red blood cell (RBC) level did not vary ($P > 0.05$). This study shows that the toad venom reduced the parasitemia in the group that was treated with 217 mg/kg toad venom which is thus comparable to the group treated with 8mg/kg standard drug (ACT). Sequel to the potency demonstrated by toad venom against plasmodial activities, it is crucial to generate a reliable and economic bufo-genic therapeutic product, while being conscious of the protection of these species. The study recommends an affirmatory research in relation to the isolation and characterization of the active anti-plasmodial agents responsible for the curative activity observed and as well as consider species-specific active compounds.

Keywords: Bufonidae; acute oral toxicity; biochemical characterization of toad venom; antiplasmodial potency of toad venom; *Plasmodium berghei*; Swiss albino mice; haematological parameters

Introduction

Malaria disease is characterized by a flu-like illness, accompanied by fever and chills; that may be fatal, if not treated [1]. It is caused by a protozoan parasite of the genus *Plasmodium* [2]. Malaria disease contributes greatly to the world disease burden and accounts for many death cases recorded in hospitals. According to a recent World malaria report released in February 2020, there were 229 million cases of malaria in 2019, up from 228 million cases in 2018 and 219 million cases in 2017 respectively [3,4]. The estimated number of malaria deaths stood at 435 000 in 2017, a similar number to the previous year. Nigeria among other African countries recorded more death cases due to malaria with about 25% malaria occurrence [1]. This is apparently an indication that malaria is highly endemic in Nigeria. This presents a severe economic and health burden to the country with about 97% of the population at risk of infection [5].

Four species of *Plasmodium* are recognized to cause malaria in man, this includes: *Plasmodium falciparum*, *P. vivax*, *P. malariae*, and *P. ovale*, transmitted by over 70 *Anopheles* species. Among the malaria parasites, *P. falciparum* is the most severe and blamable for 90% of all malaria deaths [6]. *P. berghei* have been described in African murine malaria with others being *Plasmodium chabaudi*, *Plasmodium vinckei* and *Plasmodium yoelii*. Due to *P. berghei* ability to infect rodents and the relative ease of genetic engineering, it is thus used as a well-known model organism for the investigation of malaria in man [7]. This makes this species efficient in research programs for development and screening of anti-malarial drugs and malaria vaccines that are effective [8].

Plasmodium life cycle is intricate, involving many developmental stages and sites, both in the vector and the animal host [9]. Although its intricate infectious cycle presents several sites for the development of specific drugs, finding a universal drug against malaria has been a perplexing task [10]. The main problem with the presently available anti-malarial drugs is the constant resistance developed by some parasites as opined by [4, 11-14]. The mode of action of the drugs used at the present time for the treatment of some parasitic infections are yet not very known, aiding mutations and initiating drug resistance by the parasites [15-18]. Artemisinin resistance has been described by White [12] and Dondorp et al. [19] in malaria endemic areas, threatening malaria control, treatment, and eradication of global efforts. As it is arising with anti-bacterial drugs, the abuse and the mono-therapeutic use of anti-parasitic drugs is also leading to resistance, increasing

preoccupation about a growing arrival of new cases and inefficacy on the treatment of recurrent cases [12, 19-21]. The utilization of natural products has delivered a potential strategy for identifying novel antimalarial drugs. The utilization of animals, fungi, bacteria and plants are important springs of biologically active substances with structural variety and novel mechanisms of action, which can possibly provide patentable products [22-25].

The skin secretions and venom of amphibians are rich sources of bioactive compounds, such as peptides, bufadienolides, alkaloids, proteins and biogenic amines, which have contributed meaningfully to human health and most pharmaceutical researches [26,27]. Yang et al. [27] maintained that the toad toxins from the parotoid or skin glands have significant therapeutic value for a good number of diseases. Against this backdrop, this study screened toad venoms for its antiplasmodial potency in Swiss albino mice infected with *P. berghei*.

Material and Methods

Toads Collection

Toads were collected from July to August 2019 between the hours of 0700 and 1000 from rice fields located on Latitude 8° 35'0.09" E and Longitude 8° 28'18.45" N beside the fence of the Take-Off Site of Federal University of Lafia, Nasarawa State, Nigeria. They were collected into a ventilated container and conveyed to the Department of Zoology Laboratory in the Permanent Site of Federal University of Lafia for extraction.

Extraction Of Crude Venom Extract From Toad

The process of extraction of toad venom was according to Gao et al. [28]. The toad was massaged and then the parotoid macro-glandules was pressed out. The toad was then released to its natural environment. The collected secretion was lyophilized and stored in a freezer at the Federal University of Lafia.

Biochemical Characterization

Determination of the quantitative and qualitative chemical components of the venom was achieved by Mass Spectrometry using a Mass spectrophotometer at the Spectral Laboratory and Services, Tudun wada Kaduna South, Kaduna State, Nigeria. The Shimadzu Fourier transform Infrared Spectrophotometer- FTIR 8400 S was used for the determination of the functional groups present in the venom. Gas Chromatography Analysis of toad venom was done using gas chromatography (Perkin-Elmer 8500). The Scanning Electron Microscope energy dispersive X-ray

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on *Plasmodium berghei* in Swiss Albino Mice. *Op Acc J Bio Sci & Res* 8(1)-2021.

DOI: 10.46718/JBGSR.2021.08.000190

spectroscopy (SEM-EDS) Phenom Prox, manufactured by phenom World Eindhoven (Netherlands) was used to carry out the morphology analysis (that is; analysis of the chemical elements present in the toad venom).

Experimental Animals

Twenty-five laboratory albino mice, weighing between 14g and 37g were purchased from National Veterinary Research Institute Vom, Plateau State. All animals were fed with formulated feeds and water was administered *ad libitum*. The animals were allowed to acclimatize for a period of 7 days prior to their randomization into the various experimental groups.

Coding And Weighing Of Animals

The mice were marked with a permanent ink markers on different anatomical parts of the body of each animal. These include, the head (HD), back (BK), tail (TL), right leg (RL), left leg (LL), right side (RS), left side (LS), right arm (RA), left arm (LA), back strip (BKST), head and back (HDBK), head and tail (HDTL), head and right leg (HDRL), head and left leg (HDLL), head and right side (HDRS), head and left side (HDLS), head and right arm (HDRA), head and left arm (HDLA), head and back strip (HDBKST), back and tail (BKTL), back and right leg (BKRL), back left leg (BKLL), back and right side (BKRS), back and left side (BKLS), back and right arm (BKRA). The weight of each animal was measured using the electronic compact scale.

Toxicity Study

The median oral lethal dose of the toad venom was determined in mice according to Lorke [29]. The toxicity test was carried out in two phases at the *in-vivo* laboratory of National Institute for Pharmaceutical Research and Development (NIPRID), Federal Capital Territory, Abuja, Nigeria.

Phase 1: Nine Swiss albino mice were divided into three groups with each having three individuals. The groups were administered varying doses of 10, 100 and 1000 mg/kg respectively of the toad venom and then placed under observation for 24hours in order to monitor their behaviour as well as mortality.

Phase 2: Another set of three animals, distributed into three groups of one animal each were administered varying doses of 250, 500 and 750 mg/kg respectively of the toad venom and observed for 24hours in relation to behaviour and mortality.

Then the LD_{50} was calculated by the formula by Lorke

[29]:

$$LD_{50} = \sqrt{(D_0 \times D_{100})}$$

Where

D_0 = Highest dose that gave no mortality,

D_{100} = Lowest dose that produced mortality.

Grouping, Infection And Treatment Of Infected Swiss Albino Mice

Twenty-five mice were marked and infected with *Plasmodium berghei* obtained from NIPRID and randomly divided into five groups of five mice per group. Parasites were maintained through serial blood passage in mice wherein the mice previously infected with *Plasmodium berghei* and with a high parasitemia level served as the donor. Blood samples was taken from the donor and diluted with phosphate-buffered saline such that 0.2 ml injected intraperitoneally into the experimental animals contained 1×10^7 infected erythrocytes. The blood of the mice were screened for infection before inoculation with parasite.

The curative method described by Ryley and Peters [30] was used. Treatment commenced following confirmation of parasitemia in the mice. Groups 1, 2 and 3 were treated with 217 mg/kg, 173 mg/kg and 130 mg/kg respectively of crude toad venom extract whereas group 4 mice were treated with 8 mg/kg of Artemisinin Combination Therapy (Arthemeter-lumefantrin) tablet which was crushed and diluted in distilled water before administration, while those in group 5 were infected and received no treatment (control group). Administration of treatment was done orally once daily by gavage using a mouse feeding tube for four days for the standard drug used. The parasitemia of experimental mice was establish before every treatment was administered.

Determination of Parasitemia

Blood samples were collected by bleeding the mice through the tail vein. Thin blood smears were made on clean glass microscope slides. The films were dried in air and then fixed in methanol and stained with 10% Giemsa solution [31]. The stained film was then observed under the binocular compound microscope and viewed for parasitemia. The percentage parasitemia were determined by counting the number of parasitized red blood cells (pRBC) for at least six different fields.

Percentage Parasitemia = (Parasitized RBCs)/(Total RBCs) x 100

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on Plasmodium berghei in Swiss Albino Mice. Op Acc J Bio Sci & Res 8(1)-2021.

DOI: 10.46718/JBGSR.2021.08.000190

Hematological Parameters

Using the methods described by Cheesbrough [32], the Packed Cell Volume (PCV), haemoglobin (Hb) and erythrocyte (RBC) counts were determined. These parameters were determined for each mouse before infection and after treatment. Blood samples were collected from the tail of each mouse with a heparinized capillary tube with one end sealed with plasticine.

Data Analysis

Data obtained were analyzed using R Console software (Version 3.2.2). Shapiro-wilk normality test was carried out and the data was observed to be normally distributed. One-way analysis of variance (ANOVA) was used to compare the mean parasitemia in albino mice in relation to toad venom and ACT treatments. Also, mean change in body weight and as well in haematological parameters were compared using one-way ANOVA. ANOVA test was followed by a post-hoc Tukey's Honest Significant Difference (Tukey HSD) test where there was significant difference between the means of the treatments. The level of significance was set at $P < 0.05$.

Table 1: Analysis of the chemical components of toad venom crude extract using the GC-MS.

Peak	Rate	Library/ID of compounds	Percentage of total compounds (%)
1	37.4 71	n-Hexadecanoic acid n-Hexadecanoic acid n-Hexadecanoic acid	29.04
2	38.2 84	1-Hexadecyne 5-Hexadecyne 5-Eicosyne	3.87
3	38.6 15	9,12-Octadecadienoic acid (Z,Z)- 9,12-Octadecadienoic acid (Z,Z)- 9,17-Octadecadienal, (Z)-	49.76
4	38.7 34	Octadecanoic acid Octadecanoic acid Pentadecanoic acid	7.03
5	39.3 41	Hexadecanedioic acid Dodecanoyl chloride 4-Cyclopropylme thylbenzotrile	1.87
6	40.2 35	Butyl 9,12-octadecadienoate 6-Dodecane 7,10-Hexadecadienoic acid, methyl ester	2.95
7	43.2 07	Squalene Squalene Supraene	5.48

Table 2: Protein and non-protein functional groups present in toad venom crude extract at different peaks using the FTIR.

Amino- group	Nitro- group	Non-protein group
Triethylamine	Nitro group with broad stretching of NO_2	Aromatic group (Peak 4)
Diethylamide	M-nitrotoluene	Aromatic group (Peak 14)
Aniline with a concentration of 84.653%	Nitrile group	Carbon-Hydrogen stretch with strong intensity of 89.60%
Secondary Amines	Nitrogen-Hydrogen Oxygen stretch of 93.18%	Olefins with weak band of 53.975%
n-butylamine & Benzamide with intensity (concentration) of 90.201%	Nitro-Methane with concentration of 91.31%	
Nitrogen-Hydrogen strong bond (amide) with concentration of 90.462	Nitrile group with Concentration of 89.86 %	
N-H bending		

Table 3: Analysis of chemical elements present in the toad venom crude extract using the SEM-EDS.

Element Number	Element Symbol	Element Name	Atomic Conc.	Weight Conc.
6	C	Carbon	61.76	41.26
8	O	Oxygen	25.04	22.29
25	Mn	Manganese	4.12	12.59
82	Pb	Lead	0.94	10.84
24	Cr	Chromium	1.43	4.13
7	N	Nitrogen	2.67	2.08
53	I	Iodine	0.24	1.68
11	Na	Sodium	1.12	1.44
16	S	Sulfur	0.80	1.44
9	F	Fluorine	1.28	1.36
13	Al	Aluminium	0.60	0.90

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on Plasmodium berghei in Swiss Albino Mice. Op Acc J Bio Sci & Res 8(1)-2021.

DOI: 10.46718/JBGSR.2021.08.000190




Figure 1: Mass spectroscopy of toad venom crude extract showing the abundance of the chemical compounds at different peaks.

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on Plasmodium berghei in Swiss Albino Mice. *Op Acc J Bio Sci & Res* 8(1)-2021.

DOI: 10.46718/JBGSR.2021.08.000190




Figure 2: The functional groups present in the toad venom crude extract at different peaks.




Figure 3: Mean parasitemia in Swiss albino mice in relation to treatments with toad venom and ACT.

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on Plasmodium berghei in Swiss Albino Mice. *Op Acc J Bio Sci & Res* 8(1)-2021.

DOI: 10.46718/JBGSR.2021.08.000190

Results

Determination of the Biochemical Compounds in the Toad Venom Crude Extract

The results of the biochemical analysis of the toad venom crude extract showed that 9,12-Octadecadienoic acid (Z,Z) compound was the most dominant at peak 3 as shown in Figure 1 which had a rate of 38.615 at a proportion of 49.76% (Table 1) followed by n-Hexadecanoic acid (29.04%) then Octadecanoic acid (7.03%), Squalene (5.48%), 1-Hexadecyne (3.87%), Butyl 9,12-octadecadienoate (2.95%) while Hexadecanedioic acid was the least compound (1.87%). (Figure 2) showed that the toad venom crude extract contains 18 active functional groups in which Nitrogen-Hydrogen Oxygen stretch of 93.18% was the most active functional group. The toad venom crude extract was predominantly characterized by the presence of the amino group made up of 8 functional groups, followed by the nitro-group having 6 functional groups while the non-protein group was made up of only 4 functional group as shown in Table 2.

The SEM-EDS analysis revealed that the toad venom crude extract contains 11 chemical elements, out of which carbon had the highest atomic concentration and weight of 61.76 mol/dm³ and 41.04 g/mol respectively, followed by oxygen (25.04 mol/dm³ and 22.29 g/mol) while iodine and aluminium had the least atomic concentration and weight of 0.24 mol/dm³ and 0.90g/mol respectively (Table 3).

Determination Of The Oral Toxicity (LD₅₀) Of The Toad Venom Crude Extract

The oral toxicity experiment conducted using the Lorke's method indicated that the toad crude venom extract caused mortality at a dose of 1000 mg/kg after one hour. Hyperactivity, convulsion and constant stooling were observed prior to the death of the animals. However, on the administration of 750 mg/kg, hyperactivity, diarrhea and calmness were observed within one hour, but there was no mortality recorded. The oral toxicity of the venom established was 866 mg/kg.

Mean Parasitemia In Swiss Albino Mice Across Treatments With Toad Venom And Act

The overall parasitemia was highest in the control group and least in the group treated with ACT. Therefore, the mean parasitemia in albino mice in relation to treatments showed a very high significant difference ($F_{120} = 10.34$, Adjusted $R^2 = 0.2315$, $P < 0.0001$, Figure 3). A closer look at the parasitemia in albino mice between those treated with the highest dosage of toad venom and ACT treatments showed

no significant difference ($P > 0.05$) as shown in Table 4.

Change In Body Weight Of Swiss Albino Mice After Four Days Of Treatment

After the treatments, the highest weight gain was observed in treatment with 8 mg/kg ACT (Group 5) and the least weight in the control group (infected but not treated). Thus, the mean change in body weight of albino mice after four days of parasitemia treatment in relation to treatments with toad venom and ACT showed a very high significant difference ($F_{20} = 6.694$, Adjusted $R^2 = 0.4869$, $P = 0.001374$), (Figure 4, Table 5).

Mean change in haematological conditions of albino mice after four days of treatment:

Hemoglobin (Hb)

The mean change in Hb level in albino mice after four days in relation to treatments with toad venom and ACT showed a significant difference ($F_{10} = 4.863$, Adjusted $R^2 = 0.5246$, $P = 0.01942$), (Table 6, Figure 5).

Mean Change In Red Blood Cell

The mean change in RBC level in albino mice after four days of parasitemia treatment in relation to treatments with toad venom and ACT showed no significant difference ($F_{10} = 1.929$, Adjusted $R^2 = 0.2098$, $P = 0.1821$, Table 6, Figure 6).

Mean Change In The Pcv Level

The mean change in PCV level in albino mice after four days of parasitemia treatment in relation to treatments with toad venom and ACT showed a very high significant difference ($F_{10} = 6.014$, Adjusted $R^2 = 0.5889$, $P = 0.009893$, Table 6, Figure 7).

Discussion

The biochemical analysis revealed the presence of 9,12-Octadecadienoic acid (Z,Z-) and n-Hexadecanoic acid in the toad venom which is consistent with the findings of Abubakar and Majinda [33] who recorded similar compounds from a GC-MS analysis and the preliminary antimicrobial activity of *Albizia adianthifolia* and *Pterocarpus angolensis*. Among other compounds isolated, 9,12-Octadecadienoic acid (Z,Z-) and n-Hexadecanoic acid showed the best antimicrobial activity against *E.coli*. The presence of squalene, in this study is symbolic of the antitumor potency [34] of the venom and thus, the assurance of little or no cytotoxic activity *in-vivo*. Opinions from Lozano-Grande et al. [35] suggests that squalene is a chemotherapeutic and chemo-preventive agent which inhibits tumor growth and stimulate the immune system in the treatment of diseases.

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on Plasmodium berghei in Swiss Albino Mice. Op Acc J Bio Sci & Res 8(1)-2021.

DOI: 10.46718/JBGSR.2021.08.000190

Table 4: Tukey multiple comparisons of means of parasitemia 95% family-wise confidence level.

Groups	Diff	Lower C. I.	Upper C. I.	P adj
130 mg/kg Venom-0 mg/kg Control	-0.556	-1.146 6416	0.034 64165	0.075 4076
175 mg/kg Venom-0 mg/kg Control	-0.484	-1.074 6416	0.106 64165	0.162 0644
217 mg/kg Venom-0 mg/kg Control	-1.052	-1.6426 416	-0.461 35835	0.000 0257
ACT 8 mg/kg -0 mg/kg Control	-1.216	-1.806 6416	-0.625 35835	0.000 0009
175 mg/kg Venom-130 mg/kg Venom	0.072	-0.518 6416	0.662 64165	0.997 1662
217 mg/kg Venom-130 mg/kg Venom	-0.496	-1.086 6416	0.094 64165	0.143 9029
ACT 8 mg/kg -130 mg/kg Venom	-0.660	-1.250 6416	-0.069 35835	0.020 3229
217 mg/kg Venom-175 mg/kg Venom	-0.568	-1.158 6416	0.022 64165	0.065 6025
ACT 8 mg/kg -175 mg/kg Venom	-0.732	-1.322 6416	-0.141 35835	0.007 2109
ACT 8 mg/kg -217 mg/kg Venom	-0.164	-0.754 6416	0.426 64165	0.93 89 918 ^{NS}

NS: Not Significant

Table 5: Tukey multiple comparisons of means for change in body weight between treatments 95% family-wise confidence level.

Groups	Diff	Lower C. I.	Upper C. I.	P adj
130 mg/kg Venom-0 mg/kg Control	1.8	-0.307 4476	3.907 448	0.117 6171
175 mg/kg Venom-0 mg/kg Control	1.0	-1.107 4476	3.107 448	0.622 6397
217 mg/kg Venom-0 mg/kg Control	0.8	-1.307 4476	2.907 448	0.785 9558
ACT 8 mg/kg -0 mg/kg Control	3.4	1.292 5524	5.507 448	0.0008 693*
175 mg/kg Venom-130 mg/kg Venom	-0.8	-2.907 4476	1.307 448	0.785 9558
217 mg/kg Venom-130 mg/kg Venom	-1.0	-3.107 4476	1.107 448	0.622 6397
ACT 8 mg/kg -130 mg/kg Venom	1.6	-0.507 4476	3.707 448	0.195 0802
217 mg/kg Venom-175 mg/kg Venom	-0.2	-2.307 4476	1.907 448	0.998 4504
ACT 8 mg/kg -175 mg/kg Venom	2.4	0.292 5524	4.507 448	0.020 8360*
ACT 8 mg/kg -217 mg/kg Venom	2.6	0.492 5524	4.707 448	0.011 1985*

*: Significant

Table 6: Effects of toad venom extract on the haematological parameters of Albino mice, Values were presented as Mean \pm SE; n= 5 for each group.

Treatments	Hb Count (g/dl)		RBC Count ($\times 10^9/l$)		PCV Count (%)	
	Hgb Before Treatment	Hgb After Treatment	RBC Before treatment	RBC After treatment	PCV Before Treatment	PCV After treatment
217 mg/kg Venom	10.83 \pm 0.81	11.80 \pm 1.23	6.03 \pm 1.05	5.40 \pm 0.90	40.67 \pm 0.85	42.67 \pm 0.85
173 mg/kg Venom	14.10 \pm 0.45	13.77 \pm 0.24	6.50 \pm 0.3	5.83 \pm 0.78	42.33 \pm 1.03	42.00 \pm 1.08
130 mg/kg Venom	14.03 \pm 0.31	12.57 \pm 0.65*	6.27 \pm 0.34	5.30 \pm 0.92	41.67 \pm 1.25	39.00 \pm 2.16
8 mg/kg ACT	13.97 \pm 0.06	13.80 \pm 0.25	5.83 \pm 0.39	5.23 \pm 0.66	40.33 \pm 1.03	40.33 \pm 0.62
Control (Infected & Untreated)	13.57 \pm 0.29	10.90 \pm 0.25***	6.43 \pm 0.37	3.53 \pm 0.26***	40.00 \pm 0.41	30.67 \pm 2.01***

* : $P < 0.05$

*** : $P < 0.001$




Figure 4: Mean change in body weight of albino mice in relation to treatments with toad venom and ACT after four days.

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on Plasmodium berghei in Swiss Albino Mice. Op Acc J Bio Sci & Res 8(1)-2021.

DOI: 10.46718/JBGSR.2021.08.000190




Figure 5: Mean change in hemoglobin level in albino mice in relation to treatments with toad venom and ACT after four days.




Figure 6: Mean change in RBC level in albino mice in relation to treatments with toad venom and ACT after four days.

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu IC², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on Plasmodium berghei in Swiss Albino Mice. Op Acc J Bio Sci & Res 8(1)-2021.

DOI: 10.46718/JBGSR.2021.08.000190




Figure 7: Mean change in PCV level in albino mice in relation to treatments with toad venom and ACT after four days.

Amino compounds like Triethylamine, Diethylamide and Aniline that were observed in this study agrees with the findings of Krieg et al. [36] who characterized arymethylamino steroids as a class of compounds with remarkable activity against *P. berghei* in mice and *P. falciparum* and Schistosomes in-vitro. It may be inferred that the parasite inhibition observed in this study may be as a result of the presence of these active proteins.

The presence of Carbon, Nitrogen and Oxygen is indicative that these elements make up a significant portion of the nucleic acid, amino acid, sugars and fatty acid molecules found in the toad venom [37].

The experimentation of the oral toxicity LD_{50} of the toad venom crude extract indicates that the venom becomes lethal to the laboratory albino mice at a dose of 866 mg/kg. Clinical signs of toxicity observed includes: hyperactivity, diarrhea, sedation, convulsion, constant stooling and death. The presence of lead and Chromium in the toad venom recorded in this study could have accounted for the mortality observed in the mice [38]. This agrees with Arena and Drew [39] who opined that the oral toxicity of toad venom is 0.36mg/kg and the common adult clinical dosage is 3-5mg/day, thus it becomes lethal above a dose of 135

mg/day. Although, the administration of 130 mg/kg in this study as a curative dose yielded no mortality. Similarly, this is consistent with that of Gadelha et al. [40] who observed no clinical signs of toxicity or even death on the administration of 25 mg/kg dose.

Parasitemia in Swiss albino mice decreased within the four days of treatment (Figure 3). This possibly suggests that the toad venom would be a good source of antiplasmodial drug because the parasitemia in the albino mice was almost two times less in comparison to those in control group. This concurs with the finding of Banfi et al. [41] who experimented the *in-vitro* antimalarial activities of compounds isolated from *Rhinella marina* venom (cane toad) and observed it was effective and efficient against *P. falciparum* Banfi and colleagues isolated Marinobufotoxin, Dehydrobufotenin, Marinobufagin and bufalin, out of which the Dehydrobufotenin revealed to be most active against malaria and showed the absence of cytotoxicity in-vitro. This finding is also in tandem with that of Gao et al. [28] who revealed that peptides isolated from scorpion venoms are active against *Plasmodium* species.

The low parasitemia observed in the group treated with ACT (Artemether-lumefantrine) clearly showed that it is

effective in the control of malaria parasites in animals. This is in agreement with the findings of Ihekwereme et al. [42] who reported parasite inhibition in *P. berghei* infected mice treated with the standard drug (artemether-lumefantrine). Similarly, Nyandwaro et al. [43] demonstrated the efficacy of the standard drugs in the treatment of malaria in Swiss albino.

The lack of variation in parasitemia in albino mice in relation to the highest dosage of toad venom and ACT treatments probably shows that the toad venom can serve as an alternative to the standard drug (ACT) when administered at a higher dosage since ACT was formulated primarily against *Plasmodium falciparum* in man. Correspondingly, it can be argued that chloroquine has been used for many decades against malaria in man and it is also active against murine malaria. It is therefore worthy of note that the *P. berghei* is sensitive to chloroquine [44-46]. The effective inhibition of the toad venom against *P. berghei*, *in-vivo* is in tandem with a recent finding by Banfi et al. [47] in Brazil who demonstrated an *in-vitro* inhibition of *Plasmodium falciparum* using toad venom. This is a further reflection of the isolation of peptides from animal venoms against protozoan infections, as the peptides scorpine isolated from scorpion venom was responsible for the mortality of *P. berghei* gametocytes at 15µm and 100% reduction of *P. falciparum* at 5µm [48].

Hematological parameters like hemoglobin (HBG), red blood cell (RBC) count and packed cell volume (PCV) are indicative of anemia in *Plasmodium* parasitized individuals [49]. Consequently, this research considered changes in the hematological conditions of Swiss albino mice after four days of malaria infection and associated treatments. The observed changes in hemoglobin level in albino mice after four days of parasitemia treatment in relation to treatments suggests that the toad venoms contains some anti-anemic components. This is similar with the findings Banfi et al. [41] who recorded a drastic reduction in cytotoxicity on the use of compounds isolated from the cane toads. The highest hemoglobin values were observed in the group 4 (217mg/kg venom) and the least was observed in the control group this may be as a result of the phagocytic action of the parasite on haemoglobin, consequently digesting it into haem and globulin for its consumption. This concurs with the findings of Cyril-Olutayo et al. [50] and Mokennem [51] who recorded an improved haematological parameters on the use of extracts from *Anogeissus leiocapus* and *Croton*

species respectively, from *in-vivo* studies.

The lack of variation in the mean change in RBC levels in the mice after four days of parasitemia treatment in relation to the treatment with toad venom and ACT possibly suggests that parasite targets the RBCs despite treatment. On the average, the untreated group had the least number of RBC value as a result of damage of the red cells caused by the erythrocytic stages of the malaria parasites [52,53].

The variation in the mean change in PCV level in Swiss albino mice after four days of parasitemia treatment in relation to treatment with toad venom and ACT may be as a result of the parasite inhibition pace and varying response in the experimental mice. Although the infected and untreated group had the least PCV value probably due to the destruction of red blood cells which in turn affected the percentage of the packed cells. This further shows that the reduced percentage of PCV is an indication of anaemia in malaria infected individuals. However, improved PCV values were observed in both the treatments with toad venom and ACT [54].

Conclusion

Malaria parasite success or proliferation in its host have been linked to the ability of the parasite to show resistance to host immune defense and the diverse anti-malarial drugs. However, this study has shown that pharmacological foundation of toad venom have some antiplasmodial effect against the non-human form of the parasite and could therefore be further investigated for use in the development of prospective drug for the treatment of malaria. From this findings, it is observed that the effectiveness of toad venoms at a very high dose (as was used in this study) was comparable to that of ACT. Hence, it supports the use of toad venoms in folk medicine as it is practiced by the Chinese.

Regardless of the therapeutic contributions of the bufogenic products in medicine and Public health, there is still a limited research and manufacture of toad-derived drugs. This study therefore recommends the exploration of these potent chemicals in the treatment of parasitic ailments following appropriate quantification and modification by pharmaceutical companies. Also, an affirmatory research, both *in-vivo* and *in-vitro* proceeded by the isolation and characterization of the active anti-plasmodial agents responsible for the curative activity observed and considering species-specific active compounds should be carried out.

References

- Center for Disease Control and Prevention (2017) Parasites-Malaria.
- Crutcher JM, Hoffman SL (1996) Medical Microbiology; Malaria. 4th edition. Galveston (TX): University of Texas Medical Branch at Galveston.
- World Health organization (2019a) World Malaria Report 2020.
- World Health organization (2020a). World Malaria Report 2020.
- Dawaki S, Al-Mekhlafi HM, Ithoi I, Ibrahim J, Atroosh WM, et al. (2016) Is Nigeria winning the battle against malaria? Prevalence, risk factors and KAP assessment among Hausa communities in Kano State. *Malaria Journal* 8(15): 351.
- Grimberg BT, Mehlotra RK (2011) Expanding the antimalarial drug arsenal-now but how? *Pharmaceuticals* 4: 681-712.
- Franke-Fayard B, Fonager J, Braks JA, Khan SM, Janse CJ (2010) Sequestration and Tissue Accumulation of Human Malaria Parasite: Can We Learn Anything from Rodent Models of Malaria? *PLoS Pathogens* 6(9): e1001032.
- Khan SM, Janse CJ, Kappe SH, Mikolajczak SA (2012) Genetic engineering of attenuated malaria parasites for vaccination. *Current Opinions in Biotechnology* 23(6): 908-916.
- Aly AS, Vaughan AM, Kappe SH (2009) Malaria parasite development in the mosquito and infection of the mammalian host. *Annual review of microbiology* 63: 195-221.
- Tse EG, Korsik M, Todd MH (2019) The past, present and future of anti-malarial medicines. *Malaria Journal* 18: 93.
- Bioland PB (2001) Drug Resistance in Malarial. Geneva: World Health Organization. 32.
- White NJ (2004) Antimalarial drug resistance. *Journal of Clinical Investments* 113: 1084-1092.
- Carter V, Hurd H (2010) Choosing anti-Plasmodium molecules for genetically modifying Mosquitoes: focus on peptides. *Trends in Parasitology* 26: 582-590.
- Torrent M, Pulido D, Rivas L, Andreu D (2012) Antimicrobial peptide action on Parasites. *Current Drug Targets* 13: 1138-1147.
- Teixeira C, Vale N, Pérez B, Gomes A, Gomes JRB, et al. (2014) "Recycling" Classical drugs for malaria. *Chemical Revolution* 114: 11164-11220.
- Vale N, Aguiar L, Gomes P (2014) Antimicrobial peptides: a new class of antimalarial Drugs. *Frontiers in Pharmacology* 5: 275.
- Visser BJ, Van VM, Grobusch MP (2014) Malaria: an update on current Chemotherapy. *Expert Opinion on Pharmacotherapy* 15: 2219-2254.
- Capela R, Moreira R, Lopes F (2019) An Overview of Drug Resistance in Protozoal Diseases. *Molecular Journal of International Science* 20: 574-578.
- Dondorp AM, Nosten F, Yi P, Das D, Phyto AP et al. (2009) Artemisinin Resistance in *Plasmodium falciparum* malaria. *New England Journal of Medicine* 361: 455-467.
- Chakravarty J, Sundar S (2010) Drug resistance in Leishmaniasis. *Journal of Global Infectious Diseases* 2: 167-176.
- World Health Organization (2020b) WHO status reports on artemisinin resistance and ACT efficacy.
- Rocha AB, Lopes RM, Schwartzmann G (2001) Natural products in anticancer therapy. *Current Opinion in Pharmacology* 1: 364-369.
- Vieira JGM, Dutra LA, Ferreira PMP, Moraes MO, Costa Lotufo LV, et al. (2011) Cytotoxic clerodane diterpenes from *Casearia rupestris*. *Journal of Natural Products* 74: 776-781.
- Lacerda AF, Pelegrini PB, de Oliveira DM, Vasconcelos EAR, Grossi-de-Sa MF (2016) Anti-parasitic Peptides from Arthropods and their Application in Drug Therapy. *Frontiers in Microbiology* 7:91.
- Campbell S, Soman-Faulkner K (2020) *Antiparasitic Drugs*. Treasure Island (FL): StatPearls.
- Fei L, Ye CY, Jiang JP (2003) A new bufonid genus *parapelophryne* from China (Amphibia, Anura). *Acta Zootaxonomica Sinica* 28:762-766.
- Yang Q, Zhou X, Zhang M, Bi L, Miao S, et al. (2015) Angel of human health: Current research updates in toad medicine. *American Journal of Trans. Resolutions* 7(1): 1-14.
- Gao H, Zehl M, Leitner A, Wu X, Wang Z, et al. (2010) Comparison of toad venoms from different *bufo* species by HPLC and LC-DAD-MS/MS. *Journal of ethnopharmacology* 131(2): 368-7.
- Lorke D (1983) A new approach to practical acute toxicity testing. *Archives of Toxicology* 54: 275-287.
- Ryley JF, Peters W (1970) The antimalarial activity of some quinolone esters. *Annual Tropical Medical Parasitology* 84: 209-222.
- World Health Organization (2015) Research Malaria Microscopy Standards Working Group: Microscopy for the detection, identification and quantification of malaria parasites on stained thick and thin films. Geneva: World Health Organization.
- Cheesbrough M (2004) *District Laboratory Practice in tropical Countries*. Part 2 University Press Cambridge United Kingdom 266-342.
- Abubakar MN, Majinda RT (2016) GC-MS Analysis and Preliminary Antimicrobial Activity of *Albizia adianthifolia* (Schumach) and *Pterocarpus angolensis* (DC). *Medicines* 3 (3).
- Huang Z, Lin Y, Fang J (2009) Biological and Pharmacological Activities of Squalene and Related Compounds: Potential Uses in Cosmetic Dermatology. *Molecules* 14: 540-554.
- Lozano-Grande MA, Gorinstein S, Espitia-Rangel E, Davila-Ortiz G, Martinez-Ayala AL (2018) Plant Sources; extraction methods, and uses of squalene. *International Journal of Agronomy*.
- Kriegl R, Jortzik E, Goetz A, Blandin S, Wittlin S, et al. (2017) Arylmethylamino steroids as antiparasitic agents. *Nature Communications* 8: 144-178.
- Bagrov AY, Shapiro JI, Fedora OF (2009) Endogenous cardiotonic steroids: physiology, pharmacology, and novel therapeutic targets. *Pharmacological Reviews* 61: 9-38.
- World Health Organization (2019b) Fact Sheets; Lead poisoning and health.
- Arena JM, Drew RH (1985). *Poisoning: Toxicology, Symptoms Treatments*. In: Tomas CC, editors. 5th. IL: Spring field pp. 562-564.
- Gadelha IC, de Lima JM, Batista JS, Melo MM, Soto-Blanco B (2014) Toxicity effects of toad (*Rhinella jimi* Stevaux, 2002)

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on *Plasmodium berghei* in Swiss Albino Mice. *Op Acc J Bio Sci & Res* 8(1)-2021.

DOI: 10.46718/JBGS.R.2021.08.000190

- venom in chicken *Gallus gallus domesticus*. The Scientific World Journal 2014: 8514-8573.
41. Banfi FF, Krombauer GC, da Fonseca AL, Nunes RR, Andrade SN (2021) Dehydrobufotenin extracted from the Amazonian toad *Rhinella marina* (Anura: Bufonidae) as a prototype molecule for the development of antiplasmodial drugs. *Journal of Venomous Animals and Toxins including Tropical Diseases* 27: e20200073.
 42. Ihekwereme CP, Agbata CA, Chukwueze KO, Agu, SC (2016) In vivo evaluation of antiplasmodial activity of hydroethanolic stem extract of *Baphia pubescens* in *Plasmodium berghei* infected albino mice. *Journal of Herbal Medicine and Pharmacology* 5(4): 149-152.
 43. Nyandwaro K, Oyweri J, Kimani F, Mbugua A (2020) Evaluating Antiplasmodial and Antimalarial Activities of Soybean (*Glycine max*) Seed Extracts on *P. falciparum* Parasite Cultures and *P. berghei*-Infected Mice. *Journal of Pathogens* 2020: 8
 44. Okokon JE, Ofodum KC, Ajibesin KK, Danladi B, Gamaniel KS (2005) Pharmacological Screening and evaluation of the antiplasmodial activity of *Croton zambesicus* against *Plasmodium berghei* infection in Mice. *Indian Journal of Pharmacology* 37(4): 243-246.
 45. Adetutu A, Olubukola SO, Abiodun OO, Peter A (2016) Inhibition of In Vivo Growth of *Plasmodium berghei* by *Launaea taraxacifolia* and *Amaranthus viridis* in Mice. *Malaria Research and Treatment* 2016: 9.
 46. Birru EM, Geta M, Gurmu AE (2017) Antiplasmodial Activity of *Indigofera spicata* root Extract against *Plasmodium berghei* Infection in Mice. *Malaria Journal* 16: 198.
 47. Banfi FF, Guedes KS, Andregretti CR, Anguiar AC, Debiassi BW (2016) Antiplasmodial and Cytotoxic Activities of Toad Venoms from Southern Amazon, Brazil. *Korean Journal of Parasitology* 54(4): 415-421.
 48. Carballar-Lejarazu R, Rodriguez MH, Hernnedez-Hernnedez FC, Ramos-Castaneda J, Possanic LD et al. (2008) Recombinant scorpine: multifunctional antimicrobial peptides with activity against different pathogens. *Cell Molecular Life Science* 65: 3081-3092.
 49. Al-Salahy M, Shnawa B, Abed G, Mandour A, Al-Ezzi A (2016). Parasitaemia and Its Relation to Hematological Parameters and Liver Function among Patients Malaria in Abs, Hajjah, Northwest Yemen. *Interdisciplinary Perspectives on Infectious Diseases* 2016: 5 pages.
 50. Cyril-Olutayo MC, Omonkhua AA, Akanbi OM (2013) Effects of *Anogeissus leiocarpus* on Haematological Parameters of Mice Infected with *Plasmodium berghei*. *Journal of Plant Studies* 2(2): 13-21.
 51. Mokennem LB (2015) In vivo antimalarial activity of the crude root and fruit extract of *Croton macrostachyus* against *Plasmodium berghei* in mice. *Journal of Traditional and Complementary Medicine* 5: 168-173.
 52. White NJ (2017) Malaria parasite clearance. *Malaria journal* 16(1): 88.
 53. White NJ (2018) Anaemia and malaria. *Malaria Journal* 17: 371.
 54. Etim NA, Mary E, Williams ME, Akpabio U, Offiong EEA (2017) Haematological parameters and factors affecting their values. *Agricultural Science* 2(1): 37-47.

*Corresponding author: Ayim JO, Email: joeayim2017@gmail.com

Next Submission with BGSR follows:

- Rapid Peer Review
- Reprints for Original Copy
- E-Prints Availability
- Below URL for auxiliary Submission Link: <https://biogenericpublishers.com/submit-manuscript/>

Citation: Ayim JO^{1*}, Pam VA¹, Uzoigwe NR¹, Omalu ICJ², Ombugadu A¹, Ameh SF³, Anyebe GE¹, Abe EM⁴, Tanko NS⁵, Ahmed HO¹, Aimankhu PO¹, Maikenti JI¹, Ayim TP⁶, Terna AD⁷, Okeke P⁸, Aliyu AA¹, Ayuba SO¹, Odey SA¹, Adejoh VA¹, Yina GI⁹ and Dawam, N. N.¹⁰. The Antiplasmodial Activities of Bufonidae (Toad) Venom Crude Extract on *Plasmodium berghei* in Swiss Albino Mice. *Op Acc J Bio Sci & Res* 8(1)-2021.

DOI: 10.46718/JBGSR.2021.08.000190