



Article Type: Research Article

Received: 15/08/2022

Published: 25/11/2022

DOI: 10.46718/JBGSR.2022.12.000283

Impact of COVID-19 Pandemic on Haemoparasites Prevalence

Abdel-Wahab M*, Elwethenani A, Abdullah S Al-Ghamdi, Alfahmy S, Althobaiti N, Alsofiani M, Abdulaziz H S Alkhamash, Hassan K, Bokhary M, ALfaridi M, Alkhalidy M, Alfahmi M, Alkhalidi A and Althobaiti W

¹Laboratory consultant, King Faisal Medical Complex Hospital – Taif, KSA

²Laboratory director, King Faisal Medical Complex Hospital – Taif, KSA

³Department of Laboratory, King Faisal Medical Complex Hospital – Taif, USA

⁴Laboratory specialist, King Faisal Medical Complex Hospital – Taif, KSA

⁵Laboratory specialist Al Berida pediatric hospital KSA

⁶Laboratory specialist at Administration of laboratory & Blood Bank -Taif, KSA

⁷Senior laboratory specialist Administration of laboratory & Blood Bank -Taif, KSA

⁸Nursing technician, Nursing directory – Taif, KSA

⁹Laboratory specialist, Directory of health affairs– Taif, KSA

*Corresponding author: Abdel-Wahab M, Laboratory consultant, King Faisal Medical Complex Hospital – Taif, KSA

ABSTRACT

This study has been done to propose the cause of shift regarding prevalence of haemoparasite infections. Overall, 15.2 % of samples were parasites positives; 17 % before and 8 % during the pandemic, with 60% reduction. Most of parasitic cases were in males ($p < 0.001$) showing a significant difference regarding gender between the two periods. The frequency of Infections was in patients aged 15-44 years before (15.5%) and during (8%) the Covid -19 pandemic, with significant difference observed ($p < 0.002$). Moreover, a significant difference in nationality reported, ($p = 0.024$) as the majority of patients with parasitism were non-Saudi (89.5%). Plasmodium falciparum was identified in 24 (63.2%) out of positive samples, Plasmodium vivax in 13 (34.2%), Plasmodium ovale 1 (2.6%). In coordinate with our research hypothesis, a considerable decrease in the load of haemoparasitic infections was documented with the quarantine measures taken during the Covid-19 pandemic.

KEYWORDS: Malaria; Covid -19 pandemic; Blood film

Introduction

Coronaviruses (CoV) belong to the beta-coronavirus with its high mutation rate in the family Coronaviridae causing illnesses with variable severity. Severe acute respiratory syndrome (SARS) was the first known severe one in China in 2003. A second outbreak began in 2012 in Saudi Arabia. An outbreak of a novel strain of coronavirus started on December 31 of 2019, leading to severe illness, known as SARS-CoV-2 or nCOVID-19 and killing many people with very rapid and easy spread with high infectivity rate by many routes [Song Z. et al 2019]. The WHO announced COVID-19 illness to be a global pandemic On 11 March, [Zhu, G.D. et al 2020].

Haemoparasites generally cause destruction of the red blood cells leading to anorexia, anaemia, jaundice, loss of

weight high morbidity, ending to mortality [Ademola IO. et al 2014, Opara MN. 2016].

From the fear of infections epidemics to the challenge of emerging pathogens, these mandate continuous studies for parasitic diseases detection, epidemiology description, and to devise effective case treatment and infection control [Ngui, R., et al 2012].

In this study haemoparasitic diseases were sought for in Saudi population with symptoms related to blood diseases, as two-year hospital-based studies. The objective of this study was to carry out a retrospective investigation of the haemoparasitic infections prevalence in the Saudi population from January to December period in 2019 and their corresponding 12 months in 2020.

Material and Methods

Ethical considerations: Ethical clearance was obtained from the King Faisal Medical Complex Committee (Ref: KFMC-02-T-067 to conduct this study). Informed

Consents from participants were not needed.

The study area: The investigation was conducted at KFMC in Taif at Western Saudi Arabia, one of the largest hospitals in Taif Governorate. It is a referral center for tertiary specialist care serving approximately one million residents, with a capacity of 500 beds.

Data collection: In this retrospective search, we collected data through patient medical records of Blood sample results received in the parasitology section at KFMC within the two time-periods: The first one before the pandemic (from January 2019 to December 2019) and the second was during the pandemic (from January 2020 to December 2020).

Total Samples number was 250 from hospitalized and non-hospitalized patients having fever $>38^{\circ}\text{C}$ and clinically diagnosed as malaria infection symptoms. The datasets of samples used were demographic features of patients (age, sex, and nationality) and their test results. They were tabulated, investigated, compared and endly discussion done.

The patients were 210 (84 %) males and 40 (16 %) females, with an average age of 15 - 44 years. Blood collected by vein puncture on EDTA tubes tested for malaria shortly to prevent morphological alteration of malaria parasites. Both thick and thin blood films were done , examined for the presence of Plasmodium spp. Parasites and CareStart™ Malaria RDTs were done together with the blood film which were stained with fresh 10% Giemsa's solution and examined using X 100 oil immersion magnification .If no parasites detected in 300 fields of each thick film the slides were reported negative, stained thin film preparations from positive thick films were examined for malaria species: *P. falciparum*, *Pl. vivax*, *Pl. malariae*, *Pl. ovale* or mixed infection. Finally parasitaemia was estimated in 100 fields of thin films against the leucocytes counts of the patients to check the density of infection, using the equation: number of parasites/microliter = Total parasite count/WBC count X The total leucocyte count / microliter

Data analysis: Infection prevalences in proportions and relative frequencies between group (sex and age) compared by using Chi-square test (χ^2). data analysis is conducted using SPSS version 20 (SPSS Inc, Chicago, Illinois). P-value <0.05 was considered significant.

Results

In the current study, datasets were obtained, extracted, and analysed using 250 blood samples. Table 1 describes the demographic features and patients results, both before and after the pandemic periods. Out of 200 blood samples submitted to the laboratory before the pandemic period, 175 were for male and 25 for female. Out of them , 130 were Saudi and 70 were non-Saudi residents. On the other hand, out of 50 patients samples received during the pandemic time period, 35 were males patient and 15 were females. Of them , 30 were Saudi and 20 were non-Saudi. The ages ranged between 15 and >45 years distributed into one of three groups.

Out of 250 clinically suspected cases, 38 (15.2%) were positive for Plasmodium infection using microscopy, and of this *Pl. falciparum* was present in 24(63.2%), *Pl. vivax* was in 13(34.2%), *Pl. ovale* 1(2.6%).

Before the pandemic out of 200 samples examined, 34 (17 %) were positive for one or more malaria parasites species compared to 50 specimens tested during the pandemic, 4 (8 %) were positive for one or more malaria parasite species indicating 60 % reduction in the number and proportion of parasitism between the two cohorts. In addition, the majority of positive samples were for males with a highly significant parasitism- gender relationship between the two cohorts ($p<0.001$). Before the pandemic, malaria parasites were identified in 16.5 % of male patients and in 0.5 % of female patients compared 6 % and 2 % respectively during the pandemic.

Figure 1 shows age and sex distribution of the positive patients for both pre and during the COVID-19 pandemic period. Most of parasitic infection was reported for patients aged 15-44 years both before (15.5 %) and during (8%) the pandemic compared to 1.5% & 0.0% pre & during pandemic respectively among age group >45 years.

Moreover, as shown in Figure 2, most of patients were non-Saudi residents found infected by parasites ;(89.5%), showing a significant difference in nationality reported, ($p=0.024$). Out of 70 non-Saudi patients tested, 32 (16.5 %) were found positives for malaria parasites. Compared to 2(4 %) out of 30 Saudi patients during the pandemic period (Table 1).

All malaria species, were detected more frequently in the 15-40 years age group, followed by the >40 years age group, and less <15 of age, showing highly significant difference recorded ($p<0.001$).

Table 1 shows the prevalence of malaria parasites

Table 1: Patient's demographic & descriptive malaria results, both pre – and during the pandemic

Demographic character	Pre-COVID -19 (n=200, 34 positives and 166 negative)								During COVID-19 (n= 50; 4 positives and 46 negatives)								P-value
	<15y		15-44y		>45y		Total		<15y		15-44y		>45y		Total		
	(+)	(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)	(-)	
Gender																	
Male	0	0	30	87	3	55	33	142	0	0	3	32	0	0	3	32	P<0.001
Female	0%	0%	15%	43.5%	1.5%	27.5%	16.5%	71%	0%	0%	6%	64%	0%	0%	6%	64%	
	0	0	1	14	0	10	1	24	0	0	1	9	0	5	1	14	
	0%	0%	0.5%	7%	0%	5%	0.5%	12%	0%	0%	2%	18%	0%	10%	2%	28%	
Nationality																	
Saudi	0	0	2	98	0	30	2	128	0	0	2	18	0	10	2	28	P=0.0024
Non - Saudi	0	0	29	21	3	17	32	38	0	0	2	13	0	5	2	18	
Total	0	0	31	119	3	50	34	166	0	0	4	31	0	15	4	46	
	0%	0%	15.5%	60.5%	1.5	11.5%	17%	83%	0%	0%	8%	62%	0%	30%	8%	92%	

species in the blood specimen of patients according to their ages and genders, for pre-and during the pandemic time periods.

As shown in Table 1 Before the Corona, malaria parasites were found in 16.5 % of male patients and 0.5 % of female patients. While during the Corona pandemic, male patients were positives in 6 % of samples compared to 2 % in females with highly significant difference observed between the two cohorts for all the species, (p<0.05).

As shown in Table 2 A highly significant difference observed, (p<0.001) for malaria parasite identified in samples before the pandemic.

Out of 200 clinically suspected malaria cases before the pandemic, 34(17%) were positive for Plasmodium infection out of these 22 (11 %) were *P. falciparum*, 11(5.5%) *P. vivax*, 1 (0.5 %) *P. ovale*.

On the contrary, Out of 50 clinically suspected cases during the pandemic ,4(8%) were positive for Plasmodium infection out of these 2 (4%) *P. falciparum*, 2 (4%) *P. vivax*, (0%) *P. ovale*.

Discussion

Our Present study confirm the effect of the Corona pandemic and impact of lockdown measures on the prevalence of blood parasites in Kingdom Saudi Arabia. To prove the significance of the study, results obtained for blood samples received during the pandemic were compared with those of the same time-period before the

pandemic. Taking in consideration that same months of the preceding year were chosen to do this comparison in order to avoid seasonal and climate change effects on the distribution of such parasitic agents.

Malaria is one of the most widespread infectious diseases in tropical and sub-tropical countries. There are four countries certified by The World Health Organization (WHO) as malaria free, Kingdom of Saudi Arabia is among 34 countries actively attempting to eradicate malaria. The Arabian American Oil Company (ARAMCO) in the Eastern province started malaria control in 1948, to protect employees living around this area [Meo, S.A. 2020]. A national malaria programs in 1952 was started by the Saudi Arabian government targeting malarious districts present across the kingdom planning to protect pilgrims while visiting Mecca and Medina. Transmission was arrested in the Eastern and Northern provinces in 1963 and, by the early 1970s, when Saudi Arabia joined the WHO global malaria eradication program. Malaria still persists in the provinces of Aseer and Jazan, bordering the Republic of Yemen, where a series of outbreaks the worst of them was in 1998 [Snow, R. W., et al 2013].

Our study results show that during the pandemic, a marked decrease in the total number and proportion blood parasites positive cases compared to the previous year's compared cohort. This reduction may be due to the precautions taken by the local health authority in order to decrease the spread of the Corona virus same due time may be due to the unwillingness of patients to go to hospitals




Figure 1: Age and gender distribution of parasites-positive patients before (right) and during (left) the COVID-19 pandemic time periods.

avoiding to catch the virus. Travel interdiction, washing of hands daily, and frequent use of disinfectants also may play a role in helping the reduction of infections catching.

The malaria parasites were found in 15.2 % of all specimens submitted during the two periods in our present study. In consistent with this result, Community-based studies done in various geographical areas in Saudi Arabia, have recorded higher prevalence estimates malaria parasite infection which is a major public health problem (Al Shammari et al., 2001; Al-Mohammed et al., 2010).

These higher rates can be explained by lack of hygiene and presence of unsuitable agricultural backgrounds in these areas.

Furthermore, in the present research, a highly significant patient-gender relationship between the 2 cohorts was noticed, where most of parasitic infection was observed in males, this was in coincidence with the study done by (Al Shammari et al., 2001). In the current study, males were more predisposed to infection compared to females. This disparity may be explained by behavior and




Figure 2: Distribution of age and nationality of parasites-positive patients before (right) and during (left) the COVID-19 pandemic time periods.

culture variability between the both them in Saudi Arabia.

The sex distribution in this study demonstrated that males are more affected with malaria than females in figure 1, this shows agreement with the study done by Ismail et al. [Ismail KA.et al 2019], it can explained the presence greater occupational risk for male in contracting malaria than women being exposed to moquito bites when working in mines, fields or forests at biting peak times, or migration to places where there is high endemicity, adding to this leisure activities, also sleeping habits may affect malaria

transmission [Ismail KA.et al 2019]. Also, study done by Cotter et al. [Cotter c.et al2013] prove that men may be more exposed while women may be more committed than men to apply malaria-prevention measures such as insecticide-treated bed nets.

As shown in table 3 our study showed *Pl. falciparum* 63.2% of diagnosed cases with malaria this is in contrast to result reached by Abdel-Wahab et al. [Maha M. et al 2012] as most of our patients were from Africa where *Pl. falciparum* is endemic noting that resulted cases in this

Table2: Descriptive results of Malaria pre - & during pandemic.

Applied test	Direct microscopy			P-value
	+ve (%)	-ve (%)	Total (%)	
Prepandemic	34(13.6%)	166 (66.4 %)	200 (80%)	P < 0.001
During pandemic	4 (1.6%)	46 (18.4 %)	50 (20%)	
Total	38 (15.2%)	212 (84.8%)	250 (100%)	

Table 3: Results of microscopy for detection of malaria species.

Direct microscopy	Malaria species				Total
	<i>P. vivax</i>	<i>P.falciparum</i>	<i>P.malariae</i>	<i>P. ovale</i>	
Prepandemic	11 (28.9%)	22 (57.9%)	0 (0%)	1 (2.6%)	34 (89.5%)
During pandemic	2 (5.3%)	2 (5.3%)	0 (0%)	0 (0%)	4 (10.5%)
Total	13 (34.2%)	24 (63.2%)	0 (0%)	1 (2.6%)	38 (100%)

study where non- Saudi and may be presented as relapse of previous malaria infection.

The present study was not free of constraints. The reason is that our study is considered the first study focused on investigation of blood parasite infections in humans during the Covid-19 pandemic, so it was strenuous for us to compare our data with those of others. In addition that our study was relied on medical data depending on clinical symptoms and the socio-demographic differences, while patient's residence, education, drinking, water supply and waste water disposal, work, income were not available in patients records. Such data, could provide an explanation about the distribution of blood parasites throughout the population. So the true prevalence in the country can't be estimated by hospital based study findings. Meanwhile our study depends on routine diagnostic methods in hospital having a low sensitivity for parasite detection that may under- estimate the true prevalence of parasites in addition to many other factors.

Conclusion

Infections with blood parasites remain an important problem for the studied population. A significant decrease in the distribution of these parasites was detected during the COVID-19 pandemic time period, in comparison to the corresponding pre-pandemic one. Measures of lock-down taken by the Saudi government to repress the pandemic could have a role helping this reduction. Invaluable data provided by this study will be needed for meaningful public control programs targeting reduction of the prevalence and morbidity of parasitic infections in population study.

Conflict of Interests

The authors declare that they have no conflict of interests.

Acknowledgements: The authors would like to acknowledge the Director of KFMC and KFMC Research Center for their kind cooperation.

References

- Ademola IO, Onyiche TE. Haemoparasites and haematological parameters of slaughtered Ruminants and pigs at Bodija Abattoir, Ibadan, Nigeria. Afr J Biomed Res. 2014;16(2):101–5.
- Al-Mohammed HI, Amin TT, Aboulmagd E, Hablus HR, Zaza BO. Prevalence of intestinal parasitic infections and its relationship with socio-demographics and hygienic habits among male primary schoolchildren in Al-Ahsa, Saudi Arabia. Asian Pacific Journal of Tropical Medicine, 2010; 3: 906-912. [https://doi.org/10.1016/S1995-7645\(10\)60218-0](https://doi.org/10.1016/S1995-7645(10)60218-0)
- Al Shammari S, Khoja T, El Khwasky F, Gad A. Intestinal parasitic diseases in Riyadh, Saudi Arabia: prevalence, sociodemographic and environmental associations. Tropical Medicine & International Health, 2001; 6: 184-189. <https://doi.org/10.1046/j.1365-3156.2001.00698.x>
- Cotter C, Sturrock HJ, Hsiang MS, Liu J, Phillips AA, Hwang J, et al. The changing epidemiology of malaria elimination: New strategies for new challenges. The Lancet, 2013; 382(9895): 900–911.
- Ismail KA, Gharib AF, Farghly AH, Khalifa OM. Detection of Plasmodium Antigen in Feverish Patients. Int J Hematol Blood Res, 2019; 1(1): 1-4.
- Maha M Abdel-Wahab, Khadiga A Ismail, Nagwa M El-Sayed. Laboratory Diagnosis of Malaria Infection in Clinically Suspected Cases Using Microscopic Examination, OptiMAL Rapid Antigen Test and PCR. PUJ. 2012; 5(1): 59-66.

Citation: Abdel-Wahab M*, Elwethenani A, Abdullah S Al-Ghamdi, Alfahmy S, Althobaiti N, Alsoufi M, Abdulaziz H S Alkhamash, Hassan K, Bokhary M, Alfaridi M, Alkhalidi M, Alfahmi M, Alkhalidi A and Althobaiti W. Impact of COVID-19 Pandemic on Haemoparasites Prevalence. Op Acc J Bio Sci & Res 12(1)-2022.

DOI: 10.46718/JBGSR.2022.12.000283

-
7. Meo SA. COVID-19 Pandemic: Saudi Arabia's role at national and international levels. *Journal of Diabetes Science and Technology*, 2020; 14: 758-759. <https://doi.org/10.1177/1932296820930068>
 8. Ngui R, Lim YAL, Kin LC, Chuen CS, Jaffar S. Association between anaemia, iron deficiency anaemia, neglected parasitic infections and socioeconomic factors in rural children of West Malaysia. *PLoS Neglected Tropical Diseases*, 2012; 6: e1550. <https://doi.org/10.1371/journal.pntd.0001550>
 9. Opara MN, Santali A, Mohammed BR, Jegede OC. Prevalence of haemoparasites of small ruminants in lafia nassarawa state: a guinea savannah zone of Nigeria. *J Vet Adv*, 2016; 6(6): 1251–1257.
 10. Snow RW, Amratia P, Zamani G, Mundia CW, Noor AM, Memish ZA, et al. The malaria transition on the Arabian Peninsula: progress toward a malaria-free region between 1960-2010. *Advances in parasitology*, 2013; 82: 205–251.
 11. Song Z, Xu Y, Bao L, Zhang L, Yu P, Qu Y, et al. From SARS to MERS, Thrusting Coronaviruses into the Spotlight. *Viruses*. 2019; 11
 12. Zhu GD, Cao J. Challenges and countermeasures on Chinese malaria elimination programme during the coronavirus disease 2019 (COVID-19) outbreak. *Chinese journal of schistosomiasis control*, 2020; 32: 7-9.